

# Particle-kernel approximation of the sequence of filtering density functions in state-space Markov models

Dan Crisan\*

Joaquín Míguez†

April 2, 2012

## Abstract

Sequential Monte Carlo (SMC) methods, also known as particle filters, are simulation-based recursive algorithms for the approximation of the *a posteriori* probability measures generated by state-space dynamical models. At any given time  $t$ , a particle filter produces a set of samples over the state space of the system of interest. These samples are referred to as “particles” and can be used to build a discrete approximation of the *a posteriori* probability distribution of the state, conditional on a sequence of available observations, at time  $t$ . When new observations are collected, a recursive stochastic procedure allows to update the set of particles to obtain an approximation of the new posterior at time  $t + 1$ .

One potential use of this particle set is to approximate the density associated to the *a posteriori* distribution. While practitioners have rather freely applied such density approximations in the past, the issue has received less attention from a theoretical perspective. In this paper, we address the problem of constructing kernel-based estimates of the filtering probability density function. Kernel methods are the most widely employed techniques for nonparametric density estimation and it seems natural to analyze their performance when applied to the approximate samples produced by particle filters. Here, we show how to obtain asymptotic convergence results for the particle-kernel approximations of the filtering density and its derivatives. In particular, we find convergence rates for the approximation error that hold uniformly on the state space and guarantee that the error vanishes almost surely as the number of particles in the filter grows. Based on this uniform convergence result, we first show how to build continuous measures that converge almost surely (with known rate) toward the filtering measure and then address a few applications. The latter include maximum *a posteriori* estimation of the system state using the approximate derivatives of the posterior density and the approximation of functionals of it, e.g., Shannon’s entropy.

---

\*Department of Mathematics, Imperial College London (UK). E-mail: [d.crisan@imperial.ac.uk](mailto:d.crisan@imperial.ac.uk).

†Dept. of Signal Theory and Communications, Universidad Carlos III de Madrid (Spain). E-mail: [joaquin.miguez@uc3m.es](mailto:joaquin.miguez@uc3m.es).

# 1 Introduction

## 1.1 Background

Consider two random sequences,  $\{X_t\}_{t \geq 0}$  and  $\{Y_t\}_{t \geq 1}$ , possibly multidimensional, where  $X_t$  represents the unobserved state of a system of interest and  $Y_t$  is a related observation. The aim of stochastic filtering methods is to perform inference on the state  $X_t$  given the observations  $Y_1, \dots, Y_t$ . Very often, the specific relationship between the two sequences is given by a Markov state-space model that describes

- the conditional probability distribution of  $X_t$  given  $X_{t-1}$  and
- the conditional probability distribution of the observation  $Y_t$  given the state  $X_t$ .

The probability measure that characterizes the random variable  $X_t$  conditional on the observations  $\{Y_s, 1 \leq s \leq t\}$  is usually termed the “filtering measure” and we denote it as  $\pi_t$  in the sequel. If the state-space model is linear and Gaussian,  $\pi_t$  is also Gaussian and can be computed exactly using a set of recursive equations known as the Kalman filter [32]. However, if  $X_t$  takes values in a continuous space and the model is nonlinear or non-Gaussian, the exact filter is intractable and numerical approximation techniques are necessary. The class of sequential Monte Carlo (SMC) methods, also known as particle filters [26, 33, 36, 21, 20], has become a very popular tool for this purpose. Particle filters generate discrete random measures (constructed from random samples in the state space) that can be naturally used to approximate integrals with respect to (w.r.t.) the filtering measure.

The asymptotic convergence of SMC algorithms has been well studied during the past two decades. The first formal results appeared in [12, 13], while the analysis in [8] already took into account the branching (resampling) step indispensable in most practical applications. Currently, there is a broad knowledge about the convergence of particle filters in some of the forms commonly used by practitioners; see [16, 7, 14, 35, 2, 29] and the references therein. Most of these results, however, are aimed to show that integrals of real functions w.r.t.  $\pi_t$  can be accurately approximated by weighted sums when the particle filter is run with a sufficiently large number of random samples (commonly referred to as particles). More recently, other types of convergence have been investigated. For instance, the convergence of particle approximations of maximum a posteriori (MAP) estimates of sequences has also been proved. Convergence in probability can be shown using random genealogical trees (see [41] and [14]) while almost sure convergence can also be guaranteed by extending the analysis in [9] (see [39]).

In most cases of interest, the filtering measure has a density, denoted  $p_t$ , w.r.t. a dominating measure (usually Lebesgue’s) and practitioners have freely used various estimators of this function. Less attention has been devoted to this problem from a theoretical perspective, though. Note that the samples generated by the particle filter are not drawn directly from  $p_t$ : they can only be considered as *approximate samples*, in the sense that they can be used to estimate the value of integrals with respect to the measure  $\pi_t$ . As a consequence, the convergence of a kernel density estimate of  $p_t$  built from the output of a particle filter

cannot be justified directly using the classical theory of kernel density estimation, which is concerned with samples drawn directly from the distribution of interest (see, e.g., [17, 42, 44, 43]).

The estimation of  $p_t$  is of interest by itself, since it naturally enables the computation of confidence regions, as well as MAP and maximum likelihood estimators, but also because it leads to the approximation of  $\pi_t$  by a continuous (instead of discrete) random measure. The convergence of such continuous approximations of the filtering measure in total variation distance have been investigated in the context of regularized particle filters [35] as well as for accept/reject and auxiliary particle filters [34]. In [35], two variations of the standard particle (or ‘bootstrap’ [26, 19]) filter, the pre-regularized and the post-regularized particle filters, are analyzed. The two algorithms involve the construction of kernel approximations of the filtering density, which are used to draw new particles and mitigate the sample impoverishment in the resampling step. The density estimates obtained in the pre-regularized filter are unnormalized, since the associated proportionality constants cannot be evaluated. Also, the sampling scheme in both cases is more involved than in the standard particle filters. In the pre-regularized method, a rejection sampler has to be designed (see also [40]) while in the post-regularized algorithm a mixture density has to be sampled. Under a number of regularity assumptions on the transition kernel  $\{X_t\}_{t \geq 0}$  and the conditional density of  $Y_t$  given  $X_t$ , upper bounds for the expected value of the total variation distance between  $\pi_t$  and its continuous approximations are shown in [35, Section 6].

The analysis in [34] targets the convergence of a sequence of continuous random approximations of  $\pi_t$  which are similar to those of the pre-regularized particle filter in [35]. The main difference is the use of the Markov kernel of the process  $\{X_t\}_{t \geq 0}$  for smoothing instead of a generic kernel function as in classical density estimation methods. Under adequate regularity assumptions, the convergence in probability of the total variation distance (between  $\pi_t$  and its smooth random approximations) is proved, and exponential convergence rates are obtained. The assumptions on which the analysis relies exclude some typical transition kernels (e.g., Gaussian) for the process  $\{X_t\}_{t \geq 0}$ . Also, similar to the pre-regularized particle filtering approach, the filtering density estimates can only be computed up to a proportionality constant.

## 1.2 Contributions

In this paper, we analyze the approximation of  $p_t$  constructed as the sum of properly scaled kernel functions located at the particle positions. Kernel methods are the most widely used techniques for the non-parametric estimation of probability density functions (pdf’s) and, therefore, it seems natural to analyze their convergence when applied to the approximate samples generated by particle filters.

The pdf estimators we analyze are based on generic kernel functions which are only required to satisfy mild standard conditions (essentially the same as in classical density estimation theory [42]). We describe how to build approximations in arbitrary-dimensional spaces  $\mathbb{R}^d$ ,  $d \geq 1$ , and then analyze their convergence

as the number of particles is increased and the bandwidth of the kernels is decreased. In particular, we obtain point-wise convergence rates for the absolute approximation errors, both of  $p_t$  and its derivatives<sup>1</sup> (provided they exist). The latter results can be extended to deduce uniform (instead of point-wise) convergence rates, again both for  $p_t$  and its derivatives. Specifically, we provide explicit bounds for the supremum of the approximation error and prove that it converges almost surely (a.s.) toward 0 as the number of particles is increased. Our analysis is different from the standard methods in kernel density estimation. The latter address the bias and variance of the estimators using approximations based on Taylor series (see, e.g., [42, Chapter 4] or [44, Chapter 4]) or Edgeworth expansions [27], which enable the asymptotic approximation of the mean integrated square error (MISE) of the density estimate and yield expressions involving the number of samples and the kernel bandwidth. We directly obtain convergence rates for various estimation errors (not only the MISE), given in terms of a single index that links the number of samples and the kernel bandwidth (this link is briefly discussed in Section 3.3).

The uniform (on the support of  $p_t$ ) convergence result can be exploited in a number of ways. For instance, if we let  $p_t^N$  be approximation of  $p_t$  with  $N$  particles, then we can obtain a continuous approximation of the filtering measure  $\pi_t(dx)$  as  $\tilde{\pi}_t^N(dx) = p_t^N(x)dx$ , prove that  $\tilde{\pi}_t^N$  converges to  $\pi_t$  a.s. in total variation distance (as  $N \rightarrow \infty$ ) and provide explicit convergence rates. A similar kind of analysis also leads to the calculation of convergence rates for the MISE of the particle-kernel density estimator  $p_t^N$ . Additionally, we prove that the (random) integrated square error (ISE) of a truncated version of  $p_t^N$  converges to 0 a.s. and provide convergence rates. A brief comparison of these results with the standard asymptotic approximation of the MISE for kernel estimators built from i.i.d. samples is presented at the end of Section 4.3.

The convergence in total variation distance of a continuous approximation of the filtering measure  $\pi_t$  was also addressed in [35] and [34], as mentioned in Section 1.1. Compared to these earlier contributions, our analysis guarantees the convergence of the (random) total variation distance a.s. (and with explicit rates) rather than the convergence of its expected value (as in [35]) or its convergence in probability (as in [34]). Also, our assumptions on the Markov kernel of the state process  $\{X_t\}_{t \geq 0}$  and the conditional densities of  $\{Y_t|X_t\}_{t \geq 1}$  are relatively mild and simple to check. In particular, our results also hold for light-tailed Markov kernels (e.g., Gaussian), unlike Theorems 2 and 3 in [34].

The last part of the paper is devoted to some applications of the density approximation method and the uniform convergence result. We first consider the problem of MAP estimation. We refer here to the maximization of the filtering density, a problem different from that of MAP estimation in the path space addressed, e.g., in [25, 41, 39]. We first prove that the maxima of the approximation of the filtering density

---

<sup>1</sup>Let us note here that the approximation of derivatives of the filter has received attention recently, related to problems of parameter estimation in state-space systems [11, 15]. In the latter context, the filtering pdf is made to depend explicitly on a parameter vector  $\theta = (\theta_1, \dots, \theta_d)$ , and the interest is in the computation of the partial derivatives  $\partial p_t / \partial \theta_i$  in order to implement, e.g., maximum likelihood estimation algorithms [15]. In this paper, we consider derivatives with respect to the state variables in  $X_t = (X_{1,t}, \dots, X_{d_x,t})$ , i.e.,  $\partial p_t / \partial x_{i,t}$ .

actually converge, asymptotically, to the maxima of the true function  $p_t$  and then show some simulation results that illustrate the use of gradient and Newton search algorithms on the estimated density function. Note that the latter algorithms can be “safely” used after we have proved the approximations of the first-order and second-order derivatives of  $p_t$  to be converge.

The second application we describe is the approximation of functionals of  $p_t$ . We provide first a generic result that guarantees the almost sure convergence of such approximations for bounded and Lipschitz continuous functionals. Then, we address the problem of approximating Shannon entropies [6], which is of practical interest in various machine learning and signal processing problems. The log function is neither bounded nor Lipschitz continuous and, therefore, the latter generic result does not apply to the computation of entropies. We specifically address this problem by proving first the convergence of the particle approximations of integrals of the form  $\int f(x)\pi_t(dx)$  when the test function  $f$  is unbounded. Let us remark that a large majority of the results in the literature [16, 7, 14, 35, 2] refer exclusively to the approximation of integrals of bounded functions. Only recently, the convergence of approximate integrals of unbounded test functions has been proved [30], albeit for a modified particle filter and assuming that the product of the test and the likelihood functions *is* bounded. Here, we prove the almost sure convergence of the approximations of integrals of unbounded functions for the standard particle filter, assuming only integrability of the test function. From this result, we finally deduce the almost sure convergence toward 0 of the errors in the approximation of Shannon entropies for densities with a compact support. A numerical illustration is given.

### 1.3 Organization of the paper

The rest of the paper is organized as follows. Section 2 contains background material, including a summary of notation, a description of Markov state space models and the standard particle (*bootstrap*) filter. A new lemma that establishes the convergence of the particle approximation of posterior expectations of unbounded test functions is also introduced in Section 2. The construction of particle-kernel approximations of the filtering density and its derivatives is described in Section 3. Our formal results on the convergence of the particle-kernel density estimators and the smooth approximation of the filtering measure are introduced in Section 4. This includes the point-wise and uniform approximations of  $p_t(x)$ , the convergence in total variation distance of the smooth measures  $\tilde{\pi}_t^N$  and convergence rates for the mean integrated square error and the (random) integrated square error of  $p_t^N$  and its truncated version, respectively. In Section 5 we discuss applications of the particle-kernel estimator of  $p_t$  and its derivatives. In particular, we consider the problem of (marginal) MAP estimation of the state variables and the approximation of functionals of the filtering density, including Shannon’s entropy. Finally, brief conclusions are presented in Section 6.

## 2 Particle filtering

### 2.1 Notation

We first introduce some common notations to be used through the paper, broadly classified by topics.

Below,  $\mathbb{R}$  denotes the real line, while for an integer  $d \geq 1$ ,  $\mathbb{R}^d = \overbrace{\mathbb{R} \times \dots \times \mathbb{R}}^{d \text{ times}}$

- Measures and integrals.

- $\mathcal{B}(\mathbb{R}^d)$  is the  $\sigma$ -algebra of Borel subsets of  $\mathbb{R}^d$ .
- $\mathcal{P}(\mathbb{R}^d)$  is the set of probability measures over  $\mathcal{B}(\mathbb{R}^d)$ .
- $(f, \mu) \triangleq \int f(x)\mu(dx)$  is the integral of a real function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  w.r.t. a measure  $\mu \in \mathcal{P}(\mathbb{R}^d)$ .
- Take a measure  $\mu \in \mathcal{P}(\mathbb{R}^d)$ , a Borel set  $A \in \mathcal{B}(\mathbb{R}^d)$  and a real function  $f : \mathbb{R}^d \rightarrow \mathbb{R}^+$ . The projective product  $f \star \mu$  is a measure, absolutely continuous w.r.t.  $\mu$  and proportional to  $f$ , constructed as

$$(f \star \mu)(A) = \frac{\int_A f(x)\mu(dx)}{(f, \mu)}. \quad (1)$$

- Functions.

- The supremum norm of a real function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is denoted as  $\|f\|_\infty = \sup_{x \in \mathbb{R}^d} |f(x)|$ .
- $B(\mathbb{R}^d)$  is the set of bounded real functions over  $\mathbb{R}^d$ , i.e.,  $f \in B(\mathbb{R}^d)$  if, and only if,  $\|f\|_\infty < \infty$ .
- $C_b(\mathbb{R}^d)$  is the set of continuous and bounded real functions over  $\mathbb{R}^d$ .

- Sets.

- Given a probability measure  $\mu \in \mathcal{P}(\mathbb{R}^d)$ , a Borel set  $A \in \mathcal{B}(\mathbb{R}^d)$  and the indicator function

$$I_A(x) = \begin{cases} 1, & \text{if } x \in A \\ 0, & \text{otherwise} \end{cases},$$

$\mu(A) = (I_A, \mu) = \int I_A(x)\mu(dx) = \int_A \mu(dx)$  is the probability of  $A$ .

- The Lebesgue measure of a set  $A \in \mathcal{B}(\mathbb{R}^d)$  is denoted  $\mathcal{L}(A)$ .
- For a set  $A \in \mathbb{R}^d$ ,  $A^\dagger = \mathbb{R}^d \setminus A$  denotes its complement.

- Sequences, vectors and random variables (r.v.).

- We use a subscript notation for sequences, namely  $x_{t_1:t_2} \triangleq \{x_{t_1}, \dots, x_{t_2}\}$ .
- For an element  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$  of an Euclidean space, its norm is denoted as  $\|x\| = \sqrt{x_1^2 + \dots + x_d^2}$ .
- The  $L_p$  norm of a real r.v.  $Z$ , with  $p \geq 1$ , is written as  $\|Z\|_p \triangleq E[|Z|^p]^{1/p}$ , where  $E[\cdot]$  denotes expectation.

## 2.2 Filtering in discrete-time, state-space Markov models

Consider two random sequences,  $\{X_t\}_{t \geq 0}$  and  $\{Y_t\}_{t \geq 1}$ , taking values in  $\mathbb{R}^{d_x}$  and  $\mathbb{R}^{d_y}$ , respectively. There is a common probability measure for the pair  $(\{X_t\}_{t \geq 0}, \{Y_t\}_{t \geq 1})$ , denoted  $\mathbf{P}$ , that we assume to be absolutely continuous w.r.t. the Lebesgue measure. We refer to the first sequence as the state process and we assume that it is an inhomogeneous Markov chain governed by an initial probability measure  $\tau_0 \in \mathcal{P}(\mathbb{R}^{d_x})$  and a sequence of transition kernels  $\tau_t : \mathcal{B}(\mathbb{R}^{d_x}) \times \mathbb{R}^{d_x} \rightarrow [0, 1]$ , defined as

$$\tau_t(A|x_{t-1}) \triangleq \mathbf{P}\{X_t \in A | X_{t-1} = x_{t-1}\}, \quad (2)$$

where  $A \in \mathcal{B}(\mathbb{R}^{d_x})$  is a Borel set. The sequence  $\{Y_t\}_{t \geq 1}$  is termed the observation process. Each r.v.  $Y_t$  is assumed to be conditionally independent of other observations given the state  $X_t$ , meaning that

$$\mathbf{P}\{Y_t \in A | X_{0:t} = x_{0:t}, \{Y_k = y_k\}_{k \neq t}\} = \mathbf{P}\{Y_t \in A | X_t = x_t\}, \quad (3)$$

for any  $A \in \mathcal{B}(\mathbb{R}^{d_y})$ . Additionally, we assume that every probability measure  $\gamma_t \in \mathcal{P}(\mathbb{R}^{d_y})$  in the sequence

$$\gamma_t(A|x_t) \triangleq \mathbf{P}\{Y_t \in A | X_t = x_t\}, \quad A \in \mathcal{B}(\mathbb{R}^{d_y}), \quad t = 1, 2, \dots, \quad (4)$$

has a positive density w.r.t. the Lebesgue measure. We denote this density as  $g_t(y|x)$ , hence we write  $\gamma_t(A|x_t) = \int_A g_t(y|x_t) dy$ .

The filtering problem consists in the computation of the posterior probability measure of the state  $X_t$  given a sequence of observations up to time  $t$ . Specifically, for a fixed observation record  $Y_{1:t} = y_{1:t}$ , we seek the measures  $\pi_t \in \mathcal{P}(\mathbb{R}^{d_x})$  given by

$$\pi_t(A) \triangleq \mathbf{P}\{X_t \in A | Y_{1:t} = y_{1:t}\}, \quad t = 0, 1, \dots, \quad (5)$$

where  $A \in \mathcal{B}(\mathbb{R}^{d_x})$ . For many practical problems, the interest actually lies in the computation of integrals of the form  $(f, \pi_t)$ . Note that, for  $t = 0$ , we recover the prior measure, i.e.,  $\pi_0 = \tau_0$ .

## 2.3 Particle filters

The sequence of measures  $\{\pi_t\}_{t \geq 1}$  can be numerically approximated using particle filtering. Particle filters are numerical methods based on the recursive decomposition [2]

$$\pi_t = g_t^{y_t} \star \tau_t \pi_{t-1}, \quad (6)$$

where  $g_t^{y_t} : \mathbb{R}^{d_x} \rightarrow \mathbb{R}^+$  is the function defined as  $g_t^{y_t}(x) \triangleq g_t(y_t|x)$ ,  $\star$  denotes the projective product and  $\xi_t \triangleq \tau_t \pi_{t-1}$  is the (predictive) probability measure

$$\xi_t(A) = \tau_t \pi_{t-1}(A) = \int \tau_t(A|x) \pi_{t-1}(dx), \quad A \in \mathcal{B}(\mathbb{R}^{d_x}). \quad (7)$$

Specifically, the simplest particle filter, often called ‘standard particle filter’ or ‘bootstrap filter’ [26] (see also [19]), can be described as follows.

1. **Initialization.** At time  $t = 0$  draw  $N$  i.i.d. samples from the initial distribution  $\tau_0 \equiv \pi_0$ , denoted  $x_0^{(n)}$ ,  $n = 1, \dots, N$ .
2. **Recursive step.** Let  $\Omega_{t-1}^N = \{x_{t-1}^{(n)}\}_{n=1, \dots, N}$  be the particles (samples) generated at time  $t - 1$ . At time  $t$ , proceed with the two steps below.
  - (a) For  $n = 1, \dots, N$ , draw a sample  $\bar{x}_t^{(n)}$  from the probability distribution  $\tau_t(\cdot | x_{t-1}^{(n)})$  and compute the normalized weight

$$w_t^{(n)} = \frac{g_t^{y_t}(\bar{x}_t^{(n)})}{\sum_{k=1}^N g_t^{y_t}(\bar{x}_t^{(k)})}. \quad (8)$$

- (b) For  $n = 1, \dots, N$ , let  $x_t^{(n)} = \bar{x}_t^{(k)}$  with probability  $w_t^{(k)}$ ,  $k \in \{1, \dots, N\}$ .

Step 2.(b) is referred to as resampling or selection. In the form stated here, it reduces to the so-called multinomial resampling algorithm [21, 18] but the convergence of the algorithm can be easily proved for various other schemes (see, e.g., the treatment of the resampling step in [7]). Using the samples in  $\Omega_t^N = \{x_t^{(n)}\}_{n=1, \dots, N}$ , we construct a random approximation of  $\pi_t$ , namely

$$\pi_t^N(dx_t) = \frac{1}{N} \sum_{n=1}^N \delta_{x_t^{(n)}}(dx_t), \quad (9)$$

where  $\delta_{x_t^{(n)}}$  is the delta unit-measure located at  $X_t = x_t^{(n)}$ . For any integrable function  $f$  in the state space, it is straightforward to approximate the integral  $(f, \pi_t)$  as

$$(f, \pi_t) \approx (f, \pi_t^N) = \frac{1}{N} \sum_{n=1}^N f(x_t^{(n)}). \quad (10)$$

The convergence of particle filters has been analyzed in a number of different ways. Here we only use results for the convergence of the  $L_p$  norm of the approximation errors  $(f, \pi_t^N) - (f, \pi_t)$ .

**Proposition 1** *Assume that the sequence of observations  $Y_{1:T} = y_{1:T}$  is fixed (with  $T$  being some large but finite time horizon),  $g_t^{y_t} \in B(\mathbb{R}^{d_x})$  and  $(g_t^{y_t}, \xi_t) > 0$  for every  $t = 1, 2, \dots, T$ . The following results hold.*

- (a) For all  $f \in B(\mathbb{R}^{d_x})$  and any  $p \geq 1$ ,

$$\|(f, \pi_t^N) - (f, \pi_t)\|_p \leq \frac{c_t \|f\|_\infty}{\sqrt{N}} \quad (11)$$

for  $t = 0, 1, \dots, T$ , where  $c_t$  is a constant independent of  $N$ ,  $\|f\|_\infty = \sup_{x \in \mathbb{R}^{d_x}} |f(x)|$  and the expectation is taken over all possible realizations of the random measure  $\pi_t^N$ . In particular,  $\lim_{N \rightarrow \infty} |(f, \pi_t^N) - (f, \pi_t)| = 0$  a.s. for  $0 \leq t \leq T$ .

- (b) Let  $f : \mathbb{R}^{d_x} \rightarrow \mathbb{R}$  be possibly unbounded and define  $\bar{f}_{t+1}(x) = \int f(x') \tau_{t+1}(dx' | x)$ . If  $f$  and  $\bar{f}_{t+1}$  are  $p$ -integrable (with  $p > 2$ ) w.r.t. the measure  $\pi_t$ , i.e., if  $(f^p, \pi_t) < \infty$  and  $(\bar{f}_{t+1}^p, \pi_t) < \infty$  for  $0 \leq t \leq T$ , then

$$\lim_{N \rightarrow \infty} |(f, \pi_t^N) - (f, \pi_t)| = 0 \quad \text{a.s.}$$

for  $0 \leq t \leq T$ .

See Appendix A for a proof.

**Remark 1** Part (a) of Proposition 1 is fairly standard. A similar proposition was already proved in [16], albeit under additional assumptions on the state-space model. Bounds for  $p = 2$  and  $p = 4$  can also be found in a number of references (see, e.g., [7, 10, 14]). Part (b) establishes the almost sure convergence for the approximate integrals of unbounded functions (e.g., for the approximation of the posterior mean). A similar result can be found in [30], including convergence rates. However, the analysis in [30] is carried out for a modified particle filtering algorithm, that involves a rejection test on the generated particles, and cannot be applied to the standard particle filter presented in this section.

### 3 Particle-kernel approximation of the filtering density

In the sequel we will be concerned with the family of Markov state-space models for which the posterior probability measures  $\{\pi_t\}_{t \geq 1}$  are absolutely continuous w.r.t. the Lebesgue measure and, therefore, there exist pdf's  $p_t : \mathbb{R}^{d_x} \rightarrow [0, +\infty)$ ,  $t = 1, 2, \dots$ , such that  $\pi_t(A) = \int_A p_t(x) dx$  for any  $A \in \mathcal{B}(\mathbb{R}^{d_x})$ . The density  $p_t$  is referred to as the filtering pdf at time  $t$ . In this section we briefly review the basic methodology for kernel density estimation and then describe the construction of sequences of approximations of  $p_t$  using the particles generated by a particle filter and a generic kernel function. The section concludes with the discussion on the relationship between the complexity of the particle filter (i.e., the number of particles  $N$ ) and the choice of kernel bandwidth for the density estimators.

#### 3.1 Kernel density estimators

In order to build an approximation of the function  $p_t(x)$  using a size sample of size  $N$ ,  $\{x_t^{(n)}\}_{n=1, \dots, N}$ , we resort to the classical kernel approach commonly used in density estimation [42, 44, 43]. Specifically, given a kernel function  $\phi : \mathbb{R}^{d_x} \rightarrow \mathbb{R}^+$ , we build a regularized density function of the form

$$p_t^N(x) = \frac{1}{N} \sum_{n=1}^N \phi(x - x_t^{(n)}). \quad (12)$$

In the classical theory, the kernel function  $\phi$  is often taken to be a non-negative and symmetric probability density function with zero mean and finite second order moment. Specifically, the following assumptions are commonly made [42] and we abide by them in this paper<sup>2</sup>.

**A. 1** The kernel  $\phi$  is a pdf w.r.t. the Lebesgue measure. In particular,  $\phi(x) \geq 0 \ \forall x \in \mathbb{R}^{d_x}$  and  $\int \phi(x) dx = 1$ .

---

<sup>2</sup>In classical kernel density estimation theory [42], the kernel function is often chosen to be symmetric, i.e., to satisfy  $\int x \phi(x) dx = 0$ . Such property is not needed for the results to be introduced here, though.

**A. 2** The probability distribution with density  $\phi$  has a finite second order moment, i.e.,  $c_2 = \int \|x\|^2 \phi(x) dx < \infty$ .

Given a function  $\phi$  satisfying A.1 and A.2 it is possible to define a family of re-scaled kernels

$$\phi_{\frac{1}{h}}(x) = h^{-d_x} \phi(h^{-1}x), \quad (13)$$

where  $h > 0$  is often referred to as the bandwidth of the kernel function. Both the kernel and the bandwidth can be optimized to minimize the mean integrated square error (MISE) between the regularized density and the target densities [44]. Specifically, the MISE is defined as

$$\text{MISE} \equiv \int E \left[ \left( p_t(x) - \frac{1}{N} \sum_{n=1}^N \phi_{\frac{1}{h}}(x - x_t^{(n)}) \right)^2 \right] dx, \quad (14)$$

where the expectation is taken over the random sample. Although the MISE given in Eq. (14) is intractable in general, accurate asymptotic approximations (as  $N \rightarrow \infty$ ) are known [44]. Moreover, if we assume that  $x_t^{(1)}, \dots, x_t^{(N)}$  is an i.i.d. random sample drawn exactly from  $p_t(x)$  (beware that this is *not* the case in the particle filtering framework, though), then the MISE is minimized by the Epanechnikov kernel [42]

$$\phi_E(x) = \begin{cases} \frac{d_x+2}{2v_{d_x}}(1 - \|x\|^2), & \text{if } \|x\| < 1 \\ 0, & \text{otherwise} \end{cases}, \quad (15)$$

where  $v_{d_x}$  is the volume of the unit sphere in  $\mathbb{R}^{d_x}$ . If, additionally,  $p_t(x)$  is Gaussian with unit covariance matrix, then the scaling of  $\phi_E$  that yields the minimum MISE is given by the bandwidth [44]

$$h_{opt} = [8v_{d_x}^{-1}(d_x + 4)(2\sqrt{\pi})^{d_x}]^{\frac{1}{d_x+4}} N^{-\frac{1}{d_x+4}}.$$

In our case,  $p_t(x)$  is not known (it is known not to be Gaussian in general, though) and the random sample  $x_t^{(1)}, \dots, x_t^{(N)}$  is not drawn from  $p_t(x)$ , so the standard results of [42, 44, 43] and others cannot be applied directly and a specific analysis is needed [35, 34].

In [35], two different kernel approximations of  $p_t$  were studied. Using the notation in the present paper, they can be written as

$$p_{t,pre}^N(x) \propto \frac{1}{N} \sum_{n=1}^N g_t(y_t|x) \phi_{\frac{1}{h}}(x - \bar{x}_t^{(n)}), \quad (16)$$

for the pre-regularized particle filter, and

$$p_{t,post}^N(x) = \sum_{n=1}^N w_t^{(n)} \phi_{\frac{1}{h}}(x - \bar{x}_t^{(n)}),$$

for the post-regularized particle filter. Note that  $p_{t,pre}^N(x)$  is an unnormalized approximation of  $p_t(x)$  (the normalization constant cannot be computed in general). For the post-regularized density estimator it can be shown that under certain regularity assumptions [35, Theorem 6.15]

$$E \left[ \int |p_t(x) - p_{t,post}^N(x)| dx \mid Y_{1:t} \right] \rightarrow 0 \quad \text{a.s.}$$

(where the expectation is taken w.r.t.  $p_{t,post}^N$ ) when  $N \rightarrow \infty$  and  $h \rightarrow 0$  jointly. Specifically, the mean total variation decreases as  $O(N^{-\frac{1}{2}} + h^2)$ . A similar result can be shown for  $p_{t,post}^N$  [35, Theorem 6.9].

**Remark 2** Although we use the same notation for the particles,  $\bar{x}_t^{(1)}, \dots, \bar{x}_t^{(N)}$ , as in Section 2.3, the sampling/resampling schemes in the pre-regularized and post-regularized particle filters are different from the basic ‘bootstrap’ filter [40, 35]. The pre-regularized filter, in particular, involves the use of a rejection sampler.

**Remark 3** The convergence results in [35] for the post-regularized density estimator  $p_{t,post}^N$  hold true when the following assumptions on the state-space model are guaranteed.

- The transition kernel  $R_t(x_{t-1}, A) = \int_A g_t^{Y_t}(x) \tau_t(dx|x_{t-1})$  is mixing [35, Definition 3.2].
- The likelihood satisfies that  $\sup_{u \in W^{2,1}} \frac{g_t^{Y_t} u}{\|u\|_{2,1}} < \infty$ , where  $W^{2,1}$  is the Sobolev space of functions defined on  $\mathbb{R}^{d_x}$  which, together with their derivatives up to order 2, are integrable with respect to the Lebesgue measure, and  $\|\cdot\|_{2,1}$  is the corresponding norm.
- The measure  $\tau_t(dx|x_{t-1})$  is absolutely continuous w.r.t. the Lebesgue measure, with density  $\tau_t^{x_{t-1}}(x) \in W^{2,1}$  and  $\sup_{x_{t-1} \in \mathbb{R}^{d_x}} \|\tau_t^{x_{t-1}}\|_{2,1} < \infty$ .

Assuming that  $\tau_t = \tau$  for every  $t \geq 1$  (hence, the Markov state process is homogeneous), the analysis in [34] targets the convergence in total variation distance of the continuous measure  $\rho_t^N(x)dx$ , where the density estimator  $\rho_t^N$  is defined as

$$\rho_t^N(x) = c_t \sum_{n=1}^N g_t(y_t|x) \tau^{x_{t-1}^{(n)}}(x)$$

with normalization constant  $c_t = \left( \sum_{n=1}^N \int g_t(y_t|x) \tau^{x_{t-1}^{(n)}}(x) dx \right)^{-1}$ . This is similar to the pre-regularized approximation  $p_{t,pre}^N$  but using the Markov kernel of the model,  $\tau$ , for smoothing, instead of the generic kernel  $\phi_{\frac{1}{h}}$ . Although in most problems it is possible to draw from  $\tau^{x_{t-1}}$ , it is often not possible to evaluate it and, in such cases, the approximation  $\rho_t^N$  is not practical. Also note that  $\rho_t^N$  is *not* a kernel density estimator of  $p_t$  in the classical form of Eq. (12). The sample of size  $N$  from which the approximation is constructed corresponds to the variable  $X_{t-1}$ , rather than  $X_t$ , and smoothing is achieved by way of a prediction step (using the Markov kernel  $\tau$ ). It is *not* possible, in general, to write  $\rho_t^N(x) \propto \sum_{n=1}^N g_t(y_t|x) \phi_{\frac{1}{h}}(x - x_{t-1}^{(n)})$  for some kernel function  $\phi$ . Under regularity assumptions on  $g_t$  and  $\tau$ , it is proved in [34, Theorem 2] that

$$\mathbb{P} \left\{ \int |\rho_t^N(x) - p_t(x)| dx > \epsilon \right\} \leq c_1 \exp\{-c_2 N\}, \quad t \geq 1, \quad (17)$$

for any  $\epsilon > 0$  and some constants  $c_1, c_2 > 0$ .

**Remark 4** The regularity assumptions on the state-space model in [34, Theorem 2] are the following.

(a) There are pdf's  $\{b_t\}_{t \geq 1}$  and two constants  $0 < c_\tau < C_\tau < \infty$  such that

$$c_\tau b_t(x) \leq \tau^{x_t-1}(x) \leq C_\tau b_t(x), \quad \text{for all } x, t.$$

(b) The likelihood  $g_t$  satisfies that  $\sup_{t \geq 1; x, x' \in \mathbb{R}^{d_x}; y \in \mathbb{R}^{d_y}} \frac{g_t(y|x)}{g_t(y|x')} < \infty$ .

The assumption in (a) excludes, e.g., models of the form  $X_t = h(X_{t-1}) + V_t$  where the function  $h : \mathbb{R}^{d_x} \rightarrow \mathbb{R}^{d_x}$  is not bounded or the noise process  $V_t$  is Gaussian [34, Section 4.2]. The assumption in (b) is also stronger than required for Proposition 1 to hold true.

### 3.2 Approximation of the filtering density and its derivatives

We investigate particle-kernel approximations of  $p_t$  constructed from a kernel function  $\phi$  and the samples  $x_t^{(n)}$ ,  $n = 1, \dots, N$ , generated by the particle filter. Instead of restricting our attention to procedures based on a single kernel, however, we consider a sequence of functions  $\phi_k : \mathbb{R}^{d_x} \rightarrow \mathbb{R}^+$ ,  $k \in \mathbb{N}$ , defined according to the notation in Eq. (13), i.e.,  $\phi_k(x) = k^{d_x} \phi(kx)$ . If  $\phi$  complies with A.1 and A.2, then we have similar properties for  $\phi_k$ . Trivially,  $\phi_k(x) \geq 0 \forall x \in \mathbb{R}^{d_x}$ , and it is also straightforward to check that  $\int \phi_k(x) dx = 1$ . Moreover, if we apply the change of variable  $y = kx$  and note that  $dy = k^{d_x} dx$ , then

$$\int \|x\|^2 \phi_k(x) dx = \frac{1}{k^2} \int \|y\|^2 \phi(y) dy = \frac{c_2}{k^2},$$

from A.2.

The approximation of  $p_t$  generated by the particles  $x_t^{(n)}$ ,  $n = 1, \dots, N$ , and the  $k$ -th kernel,  $\phi_k$ , is denoted as  $p_t^k$  and has the form

$$p_t^k(x) \triangleq \frac{1}{N} \sum_{n=1}^N \phi_k(x - x_t^{(n)}) = (\phi_k^x, \pi_t^N),$$

where  $\phi_k^x(x') \triangleq \phi_k(x - x')$ . Beware that, in our notation, we skip the dependence of  $p_t^k$  on the number of particles,  $N$ , for the sake of simplicity. In Section 4.1 we will assume a certain relationship between  $N$  and  $k$  that will be carried on through the rest of the paper and justifies the omission in the notation. Let us also remark that we do not construct  $p_t^k$  in order to approximate integrals w.r.t. the filtering measure (this is more efficiently achieved using Eq. (10)). Instead, we aim at applications where an explicit approximation of the density  $p_t$  is necessary. Some examples are considered in Section 5.

In order to investigate the approximation of derivatives of  $p_t$ , let us consider the multi-index  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{d_x}) \in \mathbb{N}^* \times \mathbb{N}^* \times \dots \times \mathbb{N}^*$ , where  $\mathbb{N}^* = \mathbb{N} \cup \{0\}$ , and introduce the partial derivative operator  $D^\alpha$  defined as

$$D^\alpha h \triangleq \frac{\partial^{\alpha_1} \dots \partial^{\alpha_{d_x}} h}{\partial x_1^{\alpha_1} \dots \partial x_{d_x}^{\alpha_{d_x}}}$$

for any (sufficiently differentiable) function  $h : \mathbb{R}^{d_x} \rightarrow \mathbb{R}$ . The order of the derivative  $D^\alpha h$  is denoted as  $|\alpha| = \sum_{i=1}^{d_x} \alpha_i$ . We are interested in the approximation of functions  $D^\alpha p_t(x)$  which are continuous, as explicitly given below.

**A. 3** For every  $x$  in the domain of  $p_t(x)$ ,  $D^\alpha p_t(x)$  exists and is Lipschitz continuous, i.e., there exists a constant  $c_{\alpha,t} > 0$  such that  $|D^\alpha p_t(x-z) - D^\alpha p_t(x)| \leq c_{\alpha,t} \|z\|$  for all  $x, z \in \mathbb{R}^{d_x}$ .

For the same  $\alpha$ , we also impose the following condition on the kernel  $\phi$ .

**A. 4**  $D^\alpha \phi \in C_b(\mathbb{R}^{d_x})$ , i.e.,  $D^\alpha \phi$  is a continuous and bounded function. In particular,  $\|D^\alpha \phi\|_\infty = \sup_{x \in \mathbb{R}^{d_x}} |D^\alpha \phi(x)| < \infty$ .

**Remark 5** Trivially, if  $D^\alpha \phi \in C_b(\mathbb{R}^{d_x})$  then  $D^\alpha \phi_k \in C_b(\mathbb{R}^{d_x})$  for any finite  $k$ . In particular,

$$\|D^\alpha \phi_k\|_\infty = k^{d_x+|\alpha|} \|D^\alpha \phi\|_\infty.$$

The approximation of  $D^\alpha p_t$  computed from the samples  $x_t^{(n)}$ ,  $n = 1, \dots, N$ , and the  $k$ -th kernel,  $\phi_k$ , has the form

$$D^\alpha p_t^k(x) = \frac{1}{N} \sum_{n=1}^N D^\alpha \phi_k^x(x_t^{(n)}) = (D^\alpha \phi_k^x, \pi_t^N). \quad (18)$$

### 3.3 Complexity of the particle filter and choice of kernel bandwidth

In the sequel, we will be concerned with the convergence of the sequence of approximations  $\{D^\alpha p_t^k\}_{k \geq 1}$  under the generic assumptions A.1–A.4. All the convergence results introduced in Sections 4 and 5 are given either as limits, for  $k \rightarrow \infty$ , or as error bounds that decrease with  $k$ .

Recall, however, that  $p_t^k(x) = (\phi_k^x, \pi_t^N)$ , i.e., the density estimator  $p_t^k$  depends both on the kernel bandwidth  $h = \frac{1}{k}$  and the number of particles  $N$ . A distinctive feature of the analysis in Sections 4 and 5 is that it links both indices by way of the inequality  $N \geq k^{2(d_x+|\alpha|+1)}$ , where  $|\alpha| = \sum_{i=1}^{d_x} \alpha_i$  is the order of the derivative  $D^\alpha$ . For  $\alpha = (0, \dots, 0)$ ,  $D^\alpha p_t^k = p_t^k$  and

$$N \geq k^{2(d_x+1)}. \quad (19)$$

Obviously,  $k \rightarrow \infty$  implies that  $N \rightarrow \infty$  and  $h \rightarrow 0$ .

This connection is useful to provide simple bounds for the approximation errors, but also because it yields guidance for the numerical implementation of the density estimators. In particular, for  $|\alpha| = 0$  and a fixed kernel bandwidth  $h = \frac{1}{k}$ , the inequality in (19) determines the minimum number of particles  $N$  that are needed in the particle filter in order to guarantee that convergence, at the rates given by the Theorems of Sections 4 and 5, holds. A lesser number of samples (i.e., some  $N < k^{2(d_x+1)}$ ) would result in an under-smoothed density  $p_t^k(x)$  with a bigger approximation error.

If the computational complexity of the particle filter is limited by practical considerations, then  $N$  is fixed and the error bounds to be introduced only hold when  $k \leq N^{\frac{1}{2(d_x+1)}}$  or, equivalently, when the kernel bandwidth is lower-bounded as  $h = \frac{1}{k} \geq N^{-\frac{1}{2(d_x+1)}}$ . A smaller bandwidth would, again, result in an under-smoothed approximation  $p_t^k(x)$ . On the other hand, since over-smoothing also increases the approximation error of kernel density estimators [42], it is convenient to choose the smallest possible bandwidth  $h$ . For fixed  $N$ , we should therefore select  $h = N^{-\frac{1}{2(d_x+1)}}$ .

## 4 Convergence of the approximations

Starting from Proposition 1, we prove that the kernel approximations of the filtering pdf,  $p_t^k(x)$ , and its derivatives converge a.s. for every  $x$  in the domain of  $p_t$ , both point-wise and uniformly on  $\mathbb{R}^{d_x}$ . We also prove that the smoothed approximating measure  $\check{\pi}_t^{N(k)}(dx) = p_t^k(x)dx$  converges to  $\pi_t$  in total variation distance and that the integrated square error of a sequence of truncated density estimators converges quadratically (in  $k$ ) toward 0 a.s. Explicit convergence rates for the approximations are given.

### 4.1 Almost sure convergence

In this section we obtain convergence rates for the particle-kernel approximation  $D^\alpha p_t^k(x)$  of Eq. (18). Depending on the support of the density  $p_t(x)$ , these rates may be point-wise or uniform (for all  $x$ ). In both cases, convergence is attained a.s. based on the following auxiliary result.

**Lemma 1** *Let  $\{\theta^k\}_{k \in \mathbb{N}}$  be a sequence of non-negative random variables such that, for  $p \geq 2$ ,*

$$E[(\theta^k)^p] \leq \frac{c}{k^{p-\nu}}, \quad (20)$$

*where  $c > 0$  and  $0 \leq \nu < 1$  are constant w.r.t.  $k$ . Then, there exists a non-negative and a.s. finite random variable  $U^\varepsilon$ , independent of  $k$ , such that*

$$\theta^k \leq \frac{U^\varepsilon}{k^{1-\varepsilon}}, \quad (21)$$

*where  $\frac{1+\nu}{p} < \varepsilon < 1$  is also a constant w.r.t.  $k$ .*

**Proof:** See Appendix B.

**Remark 6** *In Lemma 1, if the inequality (20) holds for all  $p \geq 2$  then the constant  $\varepsilon$  in (21) can be made arbitrarily small, i.e., we can choose  $0 < \varepsilon < 1$ .*

Using Lemma 1, it is possible to prove that  $D^\alpha p_t^k(x) \rightarrow D^\alpha p_t(x)$  a.s. and obtain explicit convergence rates. In order to establish a connection between the sequence of kernels  $\phi_k(x)$ ,  $k \in \mathbb{N}$ , and the sequence of measure approximations  $\pi_t^N$ ,  $N \in \mathbb{N}$ , we define the number of particles to be a function of the kernel index and denote it as  $N(k)$ . To be specific, for a given multi-index  $\alpha$ , we assume that  $N(k) \geq k^{2(d_x + |\alpha| + 1)}$ . In this way, all the convergence rates to be presented in this paper are primarily given in terms of the kernel index  $k$ . We first show that  $D^\alpha p_t^k \rightarrow D^\alpha p_t$  point-wise for  $x \in \mathbb{R}^{d_x}$ .

**Theorem 1** *Under assumptions A.1, A.2, A.3, A.4 and  $N(k) \geq k^{2(d_x + |\alpha| + 1)}$ , the inequality*

$$|D^\alpha p_t^k(x) - D^\alpha p_t(x)| \leq \frac{V^{x,\alpha,\varepsilon}}{k^{1-\varepsilon}} \quad (22)$$

*holds true, with  $V^{x,\alpha,\varepsilon}$  an a.s. finite, non-negative random variable and a constant  $0 < \varepsilon < 1$ . In particular,*

$$\lim_{k \rightarrow \infty} |D^\alpha p_t^k(x) - D^\alpha p_t(x)| = 0 \quad \text{a.s.} \quad (23)$$

**Proof:** Let us construct an approximation of  $p_t(x)$  using the kernel  $\phi_k$  and the true filtering measure  $\pi_t$ , namely,  $\tilde{p}_t^k(x) = (\phi_k^x, \pi_t)$ . Since  $\pi_t(dx) = p_t(x)dx$ , the approximation  $\tilde{p}_t^k$  is actually a convolution integral and can be written in two alternative ways using the commutative property, namely

$$\tilde{p}_t^k(x) = \int \phi_k(x-z)p_t(z)dz = \int \phi_k(z)p_t(x-z)dz. \quad (24)$$

Let us now consider the derivative  $D^\alpha p_t$ . If we apply the operator  $D^\alpha$  to  $\tilde{p}_t^k$  in (24) we readily obtain

$$D^\alpha \tilde{p}_t^k(x) = \int \phi_k(z)D^\alpha p_t(x-z)dz$$

and, using the latter expression, we can find an upper bound for the error  $|D^\alpha \tilde{p}_t^k(x) - D^\alpha p_t(x)|$ . In particular,

$$\begin{aligned} |D^\alpha \tilde{p}_t^k(x) - D^\alpha p_t(x)| &= \left| \int \phi_k(z)D^\alpha p_t(x-z)dz - D^\alpha p_t(x) \right| \\ &\leq \int \phi_k(z) |D^\alpha p_t(x-z) - D^\alpha p_t(x)| dz \end{aligned} \quad (25)$$

$$\leq c_{\alpha,t} \int \phi_k(z) \|z\| dz \quad (26)$$

$$\leq c_{\alpha,t} \sqrt{\int \phi_k(z) \|z\|^2 dz} \quad (27)$$

$$= c_{\alpha,t} \frac{\sqrt{c_2}}{k}, \quad (28)$$

where Eq. (25) follows from A.1 (namely,  $\phi \geq 0$ ), (26) is obtained from the Lipschitz assumption A.3, (27) follows from Jensen's inequality and, finally, the bound in (28) is obtained from assumption A.2. Note that  $c_{\alpha,t}$  and  $c_2$  are constants with respect to both  $x$  and  $k$ . As a consequence of (28),

$$\lim_{k \rightarrow \infty} D^\alpha \tilde{p}_t^k(x) = D^\alpha p_t(x).$$

Consider now the approximation, with  $N(k)$  particles,  $D^\alpha p_t^k = (D^\alpha \phi_k^x, \pi_t^{N(k)})$  of the integral  $(D^\alpha \phi_k^x, \pi_t)$ . From Proposition 1 and assumption A.4 we obtain

$$\|D^\alpha p_t^k(x) - D^\alpha \tilde{p}_t^k(x)\|_p = \|(D^\alpha \phi_k^x, \pi_t^{N(k)}) - (D^\alpha \phi_k^x, \pi_t)\|_p \leq \frac{\bar{c}_t k^{d_x+|\alpha|} \|D^\alpha \phi\|_\infty}{\sqrt{N(k)}}, \quad (29)$$

where we have used Remark 5 and the constant  $\bar{c}_t$  is independent of  $N(k)$  and  $x$ .

A straightforward application of the triangle inequality now yields

$$\|D^\alpha p_t^k(x) - D^\alpha p_t(x)\|_p \leq \|D^\alpha p_t^k(x) - D^\alpha \tilde{p}_t^k(x)\|_p + \|D^\alpha \tilde{p}_t^k(x) - D^\alpha p_t(x)\|_p. \quad (30)$$

The first term on the right-hand side of (30) can be bounded using (29), while the second term also has an upper bound given by<sup>3</sup> (28). Taking both bounds together, we arrive at

$$\|D^\alpha p_t^k(x) - D^\alpha p_t(x)\|_p \leq \frac{\bar{c}_t k^{d_x+|\alpha|} \|D^\alpha \phi\|_\infty}{\sqrt{N(k)}} + \frac{c_{\alpha,t} \sqrt{c_2}}{k} \leq \frac{\bar{c}_{\alpha,t}}{k}, \quad (31)$$

---

<sup>3</sup>Note that  $\|D^\alpha \tilde{p}_t^k(x) - D^\alpha p_t(x)\|_p = |D^\alpha \tilde{p}_t^k(x) - D^\alpha p_t(x)|$  because  $D^\alpha \tilde{p}_t^k(x)$  does not depend on  $\pi_t^{N(k)}$ .

where the second inequality follows from the assumption  $N(k) \geq k^{2(d_x+|\alpha|+1)}$  and  $\bar{c}_{\alpha,t} = \bar{c}_t \|D^\alpha \phi\|_\infty + c_{\alpha,t} \sqrt{c_{2,\alpha}}$  is a constant.

The inequality (31) immediately yields

$$E \left[ |D^\alpha p_t^k(x) - D^\alpha p_t(x)|^p \right] \leq \frac{\bar{c}_{\alpha,t}^p}{k^p} \quad (32)$$

and we can apply Lemma 1, with  $\theta^k = |D^\alpha p_t^k(x) - D^\alpha p_t(x)|$ ,  $\nu = 0$  and arbitrarily large  $p \geq 2$ , to obtain

$$|D^\alpha p_t^k(x) - D^\alpha p_t(x)| \leq \frac{V^{\alpha,x,\varepsilon}}{k^{1-\varepsilon}}, \quad (33)$$

where  $V^{\alpha,x,\varepsilon}$  is a non-negative and a.s. finite random variable and  $0 < \varepsilon < 1$  is a constant, both of them independent of  $k$ . The limit in Eq. (23) follows immediately from the inequality (33).

□

**Remark 7** The constant  $\bar{c}_{\alpha,t}$  of Eq. (31) is independent of the index  $k$  and the point  $x \in \mathbb{R}^{d_x}$ . The random variable  $V^{\alpha,x,\varepsilon}$  is also independent of the kernel index  $k$ , as explicitly given by Lemma 1. However, it may depend on the multi-index  $\alpha$  and the point  $x$  where the derivative of the density is approximated, hence the notation.

**Remark 8** For  $\alpha = (0, \dots, 0) = \mathbf{0}$ , the inequality (22) implies that that we can construct a particle approximation of  $p_t(x)$  that converges point-wise. In particular,  $D^{\mathbf{0}} p_t(x) = p_t(x)$  and  $D^{\mathbf{0}} p_t^k(x) = p_t^k(x) = (\phi_k^x, \pi_t^{N(k)})$ , hence Eq. (23) becomes

$$\lim_{k \rightarrow \infty} |p_t^k(x) - p_t(x)| = 0 \quad \text{a.s.} \quad (34)$$

for every  $x \in \mathbb{R}^{d_x}$ .

**Remark 9** The proof of Theorem 1 does not demand that the assumptions A.3, A.4 and  $N(k) \geq k^{2(d_x+|\alpha|+1)}$  hold for every possible  $\alpha$ , but only for the particular derivative we need to approximate. For instance, if we only aim to approximate  $p_t(x)$  (i.e.,  $\alpha = \mathbf{0}$ ), assumption A.2 implies that the distribution with density  $\phi$  must have a finite second order moment, assumption A.3 means that  $p_t$  must be Lipschitz, assumption A.4 implies that the basic kernel function  $\phi$  must be continuous and bounded, and it suffices that the number of particles satisfies the inequality  $N(k) \geq k^{2(d_x+1)}$ .

Most of the results to be given in the remaining of this paper are conditional on the assumptions A.1, A.2, A.3, A.4 and  $N(k) \geq k^{2(d_x+|\alpha|+1)}$ , the same as Theorem 1. However, they refer only to properties of  $p_t$  and its first order derivatives and, as a consequence, it is enough to assume that A.3 and A.4 hold true for  $\alpha = \mathbf{0}$  and  $\alpha = \mathbf{1} = (1, \dots, 1)$  alone. For the same reason, it suffices to assume  $N(k) \geq k^{2(2d_x+1)}$ .

Through the rest of the paper, we say that the “standard conditions” are satisfied when

- A.1 and A.2 hold true;

- *A.3 and A.4 hold true for, at least,  $\alpha = \mathbf{0}$  and  $\alpha = \mathbf{1}$ ; and*
- $N(k) \geq k^{2(2d_x+1)}$ .

If we restrict  $x$  to take values on a sequence of compact subsets of  $\mathbb{R}^{d_x}$ , then we can obtain a convergence rate for the error  $|p_t^k(x) - p_t(x)|$  that is uniform on  $x$ , instead of point-wise like in Theorem 1. For the following result we fix  $p \geq 2$  and consider the sequence of hypercubes

$$\mathcal{K}_k = [-M_k, +M_k] \times \dots \times [-M_k, +M_k] \subset \mathbb{R}^{d_x},$$

where  $M_k = Mk^{\frac{\beta}{d_x p}}$ , and  $M > 0$  and  $0 \leq \beta < 1$  are positive constants independent of  $k$ . Note that, for any fixed  $p$  and  $\beta > 0$ ,  $\lim_{k \rightarrow \infty} \mathcal{K}_k = \mathbb{R}^{d_x}$ .

**Theorem 2** *If the standard conditions are satisfied, then*

$$\sup_{x \in \mathcal{K}_k} |p_t^k(x) - p_t(x)| \leq \frac{U^\varepsilon}{k^{1-\varepsilon}},$$

where  $U^\varepsilon \geq 0$  is an a.s. finite random variable and  $0 < \varepsilon < 1$  is a constant, both of them independent of  $k$  and  $x$ . In particular,

$$\lim_{k \rightarrow \infty} \sup_{x \in \mathcal{K}_k} |p_t^k(x) - p_t(x)| = 0 \quad a.s.$$

**Proof:** For any  $x = (x_1, \dots, x_{d_x}) \in \mathcal{K}_k$  and a function  $f : \mathbb{R}^{d_x} \rightarrow \mathbb{R}$  continuous, bounded and differentiable,

$$f(x) - f(0) = \int_{-M_k}^{x_1} \dots \int_{-M_k}^{x_{d_x}} D^1 f(z) dz - \int_{-M_k}^0 \dots \int_{-M_k}^0 D^1 f(z) dz.$$

In particular, for  $x_i \in [-M_k, M_k]$ ,  $i = 1, \dots, d_x$ , and the assumption A.4 with  $\alpha = \mathbf{1}$ ,

$$|p_t^k(x) - p_t(x)| \leq 2 \int_{-M_k}^{M_k} \dots \int_{-M_k}^{M_k} |D^1 p_t^k(z) - D^1 p_t(z)| dz + |p_t^k(0) - p_t(0)| \quad (35)$$

and, as a consequence,

$$\sup_{x \in \mathcal{K}_k} |p_t^k(x) - p_t(x)| \leq 2A^k + |p_t^k(0) - p_t(0)|, \quad (36)$$

where

$$A^k = \int_{-M_k}^{M_k} \dots \int_{-M_k}^{M_k} |D^1 p_t^k(z) - D^1 p_t(z)| dz.$$

An application of Jensen's inequality yields, for  $p \geq 1$ ,

$$\left( \frac{1}{2^{d_x} M_k^{d_x}} A^k \right)^p \leq \frac{1}{2^{d_x} M_k^{d_x}} \int_{-M_k}^{M_k} \dots \int_{-M_k}^{M_k} |D^1 p_t^k(s_\ell(z)) - D^1 p_t(s_\ell(z))|^p dz,$$

hence

$$(A^k)^p \leq 2^{d_x(p-1)} M_k^{d_x(p-1)} \sum_{\ell=0}^{2^{d_x}-1} \int_0^{M_k} \dots \int_0^{M_k} |D^1 p_t^k(s_\ell(z)) - D^1 p_t(s_\ell(z))|^p dz. \quad (37)$$

Since

$$E \left[ |D^1 p_t^k(s_\ell(z)) - D^1 p_t(s_\ell(z))|^p \right] \leq \frac{\bar{c}_{1,t}^p}{k^p} \quad (38)$$

from inequality (31) in the proof of Theorem 1, we can combine (38) and (37) to arrive at

$$E[(A^k)^p] \leq \frac{2^{d_x p} M_k^{d_x p} \bar{c}_{\mathbf{1},t}^p}{k^p} = \frac{\bar{c}_{\mathbf{1},t}^p}{k^{p-\beta}},$$

where the equality follows from the relationship  $M_k = \frac{1}{2}k^{\frac{\beta}{d_x p}}$ . Using Lemma 1 with  $\theta_k = A^k$ ,  $p \geq 2$ ,  $\nu = \beta$  and  $c = \bar{c}_{\mathbf{1},t}^p$ , we obtain a constant  $\varepsilon_1 \in \left(\frac{1+\beta}{p}, 1\right)$  and a non-negative and a.s. finite random variable  $V^{A,\varepsilon_1}$ , both of them independent of  $k$ , such that

$$A^k \leq \frac{V^{A,\varepsilon_1}}{k^{1-\varepsilon_1}}. \quad (39)$$

Since, from Proposition 1,

$$E\left[|p_t^k(x) - p_t(x)|^p\right] \leq \frac{\bar{c}_{\mathbf{0},t}^p}{k^p},$$

we can apply Lemma 1 again, with  $\theta_k = |p_t^k(0) - p_t(0)|$ ,  $p \geq 2$ ,  $\nu = 0$  and  $c = \bar{c}_{\mathbf{0},t}^p$  to obtain that

$$|p_t^k(0) - p_t(0)| \leq \frac{V^{p_t(0),\varepsilon_2}}{k^{1-\varepsilon_2}}, \quad (40)$$

where  $\varepsilon_2 \in \left(\frac{1}{p}, 1\right)$  is a constant and  $V^{p_t(0),\varepsilon_2}$  is a non-negative and a.s. finite random variable, both of them independent of  $k$ .

If we choose  $\varepsilon = \varepsilon_1 = \varepsilon_2 \in \left(\frac{1+\beta}{p}, 1\right)$  and define  $U^\varepsilon = V^{A,\varepsilon_1} + V^{p_t(0),\varepsilon_2}$ , then the combination of Eqs. (36), (39) and (40) yields

$$\sup_{x \in \mathcal{K}_k} |p_t^k(x) - p_t(x)| \leq \frac{U^\varepsilon}{k^{1-\varepsilon}},$$

where  $U^\varepsilon$  is a.s. finite. Note that  $U^\varepsilon$  and  $\varepsilon$  are independent of  $k$ . Moreover, we can choose  $p$  as large as we wish and  $\beta > 0$  as small as needed, hence we can select  $\varepsilon \in (0, 1)$ . Therefore, the proof is complete.

□

**Remark 10** Assuming that A.3 and A.4 hold for the multi-index  $\alpha' = \alpha + \mathbf{1}$ , the argument of the proof of Theorem 2 can also be adapted to show that

$$\sup_{x \in \mathcal{K}_k} |D^\alpha p_t^k(x) - D^\alpha p_t(x)| \leq \frac{\tilde{U}^\varepsilon}{k^{1-\varepsilon}},$$

where the constant  $0 < \varepsilon < 1$  and the a.s. finite random variable  $\tilde{U}^\varepsilon \geq 0$  are independent of  $k$ .

**Remark 11** Theorem 2 also holds for a fixed compact subset  $\mathcal{K} \subset \mathbb{R}^{d_x}$  instead of the sequence  $\mathcal{K}_1, \mathcal{K}_2, \dots$ . In particular, the presented proof is still valid for a fixed hypercube  $\mathcal{K} = [-M, +M] \times \dots \times [-M, +M]$  (i.e., with  $\beta = 0$  and fixed  $M > 0$  in the definition of  $M_k$ ). Therefore,

$$\sup_{x \in \mathcal{K}} |p_t^k(x) - p_t(x)| \leq \frac{\tilde{U}^\varepsilon}{k^{1-\varepsilon}}, \quad (41)$$

where the constant  $0 < \varepsilon < 1$  and the a.s. finite random variable  $\tilde{U}^\varepsilon \geq 0$  are independent of  $k$ .

## 4.2 Convergence in total variation distance

The total variation distance (TVD) between two measures  $\mu_1, \mu_2 \in \mathcal{P}(\mathbb{R}^d)$  on the Borel  $\sigma$ -algebra  $\mathcal{B}(\mathbb{R}^d)$  is defined as

$$d_{TV}(\mu_1, \mu_2) \triangleq \sup_{A \in \mathcal{B}(\mathbb{R}^d)} |\mu_1(A) - \mu_2(A)|.$$

Correspondingly, a sequence of measures  $\mu^n \in \mathcal{P}(\mathbb{R}^d)$  converges toward  $\mu \in \mathcal{P}(\mathbb{R}^d)$  in TVD when  $\lim_{n \rightarrow \infty} d_{TV}(\mu^n, \mu) = 0$ . It can be shown that, if  $\mu^n$  and  $\mu$  have densities w.r.t. the Lebesgue measure, denoted  $q^n$  and  $q$ , respectively, then

$$d_{TV}(\mu^n, \mu) = \frac{1}{2} \int |q^n(x) - q(x)| dx$$

and, therefore, the sequence  $\mu^n$  converges to  $\mu$  in TVD if, and only if,

$$\lim_{n \rightarrow \infty} \int |q^n(x) - q(x)| dx = 0. \quad (42)$$

Consider the smooth approximating measures

$$\check{\pi}_t^{N(k)}(dx) = p_t^k(x)dx, \quad k = 1, 2, \dots$$

In this section we show that the sequence  $\check{\pi}_t^{N(k)}$  converges toward  $\pi_t$  in TVD, as  $k \rightarrow \infty$ , by proving first that  $\int |p_t^k - p_t| dx \rightarrow 0$  under the same assumptions of Theorem 2. This result is established by Theorem 3 below. The same as in the proof of Theorem 2, we consider an increasing sequence of hypercubes  $\mathcal{K}_1 \subset \dots \subset \mathcal{K}_k \subset \dots \subset \mathbb{R}^{d_x}$ , where  $\mathcal{K}_k = [-M_k, +M_k] \times \dots \times [-M_k, +M_k]$  and  $M_k = \frac{1}{2}k^{\frac{\beta}{d_x p}}$ , with constants  $0 < \beta < 1$  and  $p > 3$ . Also, recall that, for a set  $A \in \mathbb{R}^d$ ,  $A^\dagger = \mathbb{R}^d \setminus A$  denotes its complement and, given a probability measure  $\mu \in \mathcal{P}(\mathbb{R}^d)$ ,  $\mu(A) = \int_A \mu(dx)$  is the probability of  $A$ .

**Theorem 3** *If the standard conditions are satisfied and  $\pi_t(\mathcal{K}_k^\dagger) \leq \frac{b}{2}k^{-\gamma}$ , where  $b > 0$  and  $\gamma > 0$  are arbitrary but constant w.r.t.  $k$ , then*

$$\int |p_t^k(x) - p_t(x)| dx < \frac{Q^\varepsilon}{k^{\min\{1-\varepsilon, \gamma\}}},$$

where  $Q^\varepsilon > 0$  is an a.s. finite random variable and  $0 < \varepsilon < 1$  is a constant, both of them independent of  $k$ . In particular,

$$\lim_{k \rightarrow \infty} \int |p_t^k(x) - p_t(x)| dx = 0 \quad a.s.$$

and, as a consequence,

$$\lim_{k \rightarrow \infty} d_{TV}(\check{\pi}_t^{N(k)}, \pi_t) = 0 \quad a.s.$$

**Proof:** We start with a trivial decomposition of the integrated absolute error,

$$\begin{aligned} \int |p_t^k(x) - p_t(x)| dx &= \int_{\mathcal{K}_k} |p_t^k(x) - p_t(x)| dx + \int_{\mathcal{K}_k^\dagger} |p_t^k(x) - p_t(x)| dx \\ &\leq \int_{\mathcal{K}_k} |p_t^k(x) - p_t(x)| dx + 2 \int_{\mathcal{K}_k^\dagger} p_t(x) dx + \int_{\mathcal{K}_k^\dagger} (p_t^k(x) - p_t(x)) dx, \end{aligned}$$

where the equality follows from  $\mathcal{K}_k \cup \mathcal{K}_k^\dagger = \mathbb{R}^{d_x}$  and the inequality is obtained from the fact that  $p_t$  and  $p_t^k$  are non-negative. Moreover

$$\left| \int_{\mathcal{K}_k^\dagger} (p_t^k(x) - p_t(x)) dx \right| = \left| \int_{\mathcal{K}_k} (p_t(x) - p_t^k(x)) dx \right| \leq \int_{\mathcal{K}_k} |p_t^k(x) - p_t(x)| dx$$

hence

$$\int |p_t^k(x) - p_t(x)| dx \leq 2 \int_{\mathcal{K}_k} |p_t^k(x) - p_t(x)| dx + 2 \int_{\mathcal{K}_k^\dagger} p_t(x) dx \quad (43)$$

The first term in (43) can be bounded by

$$\int_{\mathcal{K}_k} |p_t^k(x) - p_t(x)| dx \leq \mathcal{L}(\mathcal{K}_k) \sup_{x \in \mathcal{K}_k} |p_t^k(x) - p_t(x)|, \quad (44)$$

where  $\mathcal{L}(\mathcal{K}_k) = (2M_k)^{d_x} = k^{\frac{\beta}{p}}$  is the Lebesgue measure of  $\mathcal{K}_k$ . From Theorem 2, the supremum in (44) can be bounded as  $\sup_{x \in \mathcal{K}_k} |p_t^k(x) - p_t(x)| \leq V^{\varepsilon_1} / k^{1-\varepsilon_1}$ , where  $V^{\varepsilon_1} \geq 0$  is an a.s. finite random variable and  $\frac{1+\beta}{p} < \varepsilon_1 < 1$  is a constant, both independent of  $k$ . Therefore, the inequality (44) can be extended to yield

$$\int_{\mathcal{K}_k} |p_t^k(x) - p_t(x)| dx \leq \frac{V^{\varepsilon_1}}{k^{1-\varepsilon_1-\frac{\beta}{p}}} = \frac{V^\varepsilon}{k^{1-\varepsilon}}, \quad (45)$$

where  $\varepsilon = \varepsilon_1 + \frac{\beta}{p}$  and  $V^\varepsilon = V^{\varepsilon_1}$ . If we choose  $\varepsilon_1 < 1 - \frac{\beta}{p}$ , then  $\varepsilon \in \left(\frac{1+2\beta}{p}, 1\right)$ . Note that, for  $\beta < 1$  and  $p > 3$ ,  $1 - \frac{\beta}{p} - \frac{1+\beta}{p} > 1 - \frac{3}{p} > 0$ , hence both  $\varepsilon_1$  and  $\varepsilon$  are well defined.

For the second integral in Eq. (43), note that  $\int_{\mathcal{K}_k^\dagger} p_t(x) dx = \pi_t(\mathcal{K}_k^\dagger)$  and, therefore, it can be bounded directly from the assumptions in the Theorem, i.e.,

$$2 \int_{\mathcal{K}_k^\dagger} p_t(x) dx \leq bk^{-\gamma}, \quad (46)$$

where  $b > 0$  and  $\gamma > 0$  are constant w.r.t.  $k$ . Putting together Eqs. (43), (45) and (46) gives us the result.

□

### 4.3 Integrated square error

A standard figure of merit for the assessment of kernel density estimators is the mean integrated square error (MISE) [42, 43]. If we assume that both  $p_t(x)$  and the kernel  $\phi(x)$  take values on a compact set  $\mathcal{K}$ , then it is relatively simple to prove that the MISE of the sequence of approximations  $D^\alpha p_t^k$  converges toward 0 quadratically with the index  $k$ . In particular, we have the following result.

**Theorem 4** Assume that A.1, A.2, A.3, A.4 and  $N(k) \geq k^{2(d_x+|\alpha|+1)}$  hold true. If both  $p_t(x)$  and the kernel  $\phi(x)$  take values on a compact set  $\mathcal{K} \subset \mathbb{R}^{d_x}$ , then

$$\text{MISE} \equiv \int_{\mathcal{K}} E \left[ (D^\alpha p_t^k(x) - D^\alpha p_t(x))^2 \right] dx \leq \frac{c_{\alpha, \mathcal{K}, t}}{k^2},$$

where  $c_{\alpha, \mathcal{K}, t} > 0$  is constant w.r.t.  $k$ .

**Proof:** Since any compact set is contained in a larger hypercube, we can choose  $\mathcal{K} = [-M, +M] \times \cdots \times [-M, +M]$  without loss of generality. Furthermore, since the assumptions of Theorem 1 are satisfied, we can recall the inequality in (32), which, selecting  $p = 2$ , yields

$$E \left[ (D^\alpha p_t^k(x) - D^\alpha p_t(x))^2 \right] \leq \frac{\bar{c}_{\alpha,t}^2}{k^2},$$

where the constant  $\bar{c}_{\alpha,t}^2$  is independent of  $k$  and  $x$ . Therefore,

$$\int_{\mathcal{K}} E \left[ (D^\alpha p_t^k(x) - D^\alpha p_t(x))^2 \right] dx \leq \frac{\bar{c}_{\alpha,t}^2}{k^2} \mathcal{L}(\mathcal{K}) \leq \frac{c_{\alpha,\mathcal{K},t}}{k^2},$$

where  $\mathcal{L}(\mathcal{K}) = (2M)^{d_x}$  is the Lebesgue measure of  $\mathcal{K}$  and  $c_{\alpha,\mathcal{K},t} = (2M)^{d_x} \bar{c}_{\alpha,t}^2$ .

□

It is also possible to establish a quadratic convergence rate (w.r.t.  $k$ ) for the integrated square error (ISE) of a sequence of truncated density approximations. In particular, consider the usual hypercubes  $\mathcal{K}_k = [-M_k, +M_k] \times \cdots \times [-M_k, +M_k]$  with  $M_k = \frac{1}{2} k^{\frac{\beta}{d_x p}}$ , for some  $p > \frac{5}{2}$  and a constant  $0 < \beta < 1$ , and define the truncated density estimators

$$p_t^{\top,k}(x) = I_{\mathcal{K}_k}(x) p_t^k(x) = \begin{cases} p_t^k(x), & \text{if } x \in \mathcal{K}_k, \\ 0, & \text{otherwise} \end{cases}.$$

Since  $\lim_{k \rightarrow \infty} \mathcal{K}_k = \mathbb{R}^{d_x}$ , it follows that  $\lim_{k \rightarrow \infty} |p_t^{\top,k}(x) - p_t^k(x)| = 0$  and we can make  $p_t^{\top,k}$  arbitrarily close to the original approximation. The theorem below states that  $p_t^{\top,k}$  converges a.s. toward  $p_t$ , with a quadratic rate.

**Theorem 5** *If the standard conditions are satisfied,  $p_t \in B(\mathbb{R}^{d_x})$  and  $\pi_t(\mathcal{K}_k) \leq b k^{-\gamma}$ , where  $b > 0$  and  $\gamma > 0$  are arbitrary but constant w.r.t.  $k$ , then*

$$ISE \equiv \int \left( p_t^{\top,k}(x) - p_t(x) \right)^2 \leq \frac{U^\varepsilon}{k^{2-\varepsilon}},$$

where  $U^\varepsilon \geq 0$  is an a.s. finite random variable, independent of  $k$ , and  $0 < \varepsilon < 2$  is an arbitrarily small constant. In particular,

$$\lim_{k \rightarrow \infty} \int \left( p_t^{\top,k}(x) - p_t(x) \right)^2 dx = 0 \quad \text{a.s.}$$

**Proof:** We start with the trivial decomposition

$$\int \left( p_t^{\top,k}(x) - p_t(x) \right)^2 dx = \int_{\mathcal{K}_k} \left( p_t^{\top,k}(x) - p_t(x) \right)^2 dx + \int_{\mathcal{K}_k^c} \left( p_t^{\top,k}(x) - p_t(x) \right)^2 dx, \quad (47)$$

where  $\mathcal{K}_k^c = \mathbb{R}^{d_x} \setminus \mathcal{K}_k$  is the complement of  $\mathcal{K}_k$ , and, expanding the square in the last integral of Eq. (47), we obtain

$$\begin{aligned} \int \left( p_t^{\top,k}(x) - p_t(x) \right)^2 dx &= \int_{\mathcal{K}_k} \left( p_t^{\top,k}(x) - p_t(x) \right)^2 dx + \int_{\mathcal{K}_k^c} \left( p_t(x) - p_t^{\top,k}(x) \right) p_t(x) dx \\ &\quad + \int_{\mathcal{K}_k^c} \left( p_t^{\top,k}(x) - p_t(x) \right) p_t^{\top,k}(x) dx. \end{aligned} \quad (48)$$

In the rest of the proof, we compute upper bounds for each of the integrals on the right-hand side of Eq. (48).

For the first term in (48) we note that  $p_t^{\top,k}(x) = p_t^k(x)$  for all  $x \in \mathcal{K}_k$ , hence

$$\int_{\mathcal{K}_k} \left( p_t^{\top,k}(x) - p_t(x) \right)^2 dx = \int_{\mathcal{K}_k} \left( p_t^k(x) - p_t(x) \right)^2 dx \leq \mathcal{L}(\mathcal{K}_k) \left( \sup_{x \in \mathcal{K}_k} \left| p_t^{\top,k}(x) - p_t(x) \right| \right)^2, \quad (49)$$

where  $\mathcal{L}(\mathcal{K}_k) = (2M_k)^{d_x} = k^{\frac{\beta}{p}}$ . Using Theorem 2, we obtain an upper bound for the supremum in Eq. (49), namely  $\sup_{x \in \mathcal{K}_k} |p_t^k(x) - p_t(x)| \leq V^{\varepsilon_1} / k^{1-\varepsilon_1}$ , where  $V^{\varepsilon_1} \geq 0$  is an a.s. finite random variable and  $\frac{1+\beta}{p} < \varepsilon_1 < 1$  is a constant. Both  $V^{\varepsilon_1}$  and  $\varepsilon_1$  are independent of  $k$ . We then extend the inequality in (49) as

$$\int_{\mathcal{K}_k} \left( p_t^{\top,k}(x) - p_t(x) \right)^2 dx \leq k^{\frac{\beta}{p}} \frac{(V^{\varepsilon_1})^2}{k^{2-2\varepsilon_1}} = \frac{\tilde{U}^\varepsilon}{k^{2-\varepsilon}}, \quad (50)$$

where  $\varepsilon = 2\varepsilon_1 + \frac{\beta}{p}$  and  $\tilde{U}^\varepsilon = (V^{\varepsilon_1})^2$ . If we choose  $\varepsilon_1 < 1 - \frac{\beta}{2p}$ , then  $\varepsilon \in \left( \frac{2+3\beta}{p}, 2 \right)$ . Note that, for  $\beta < 1$  and  $p > \frac{5}{2}$ ,  $2 - \frac{2+3\beta}{p} > 0$ , hence  $\varepsilon$  is well defined.

For the second term in the right-hand side of Eq. (48) we simply note that  $p_t^{\top,k}(x) = 0$  for all  $x \in \mathcal{K}_k^\dagger$  and  $p_t(x) < \|p_t\|_\infty < \infty$ , since  $p_t \in B(\mathbb{R}^{d_x})$ . Therefore,

$$\int_{\mathcal{K}_k^\dagger} \left( p_t(x) - p_t^{\top,k}(x) \right) p_t(x) dx \leq \|p_t\|_\infty \int_{\mathcal{K}_k^\dagger} p_t(x) dx = \|p_t\|_\infty \pi_t(\mathcal{K}_k^\dagger),$$

and using the assumption  $\pi_t(\mathcal{K}_k^\dagger) \leq bk^{-\gamma}$  we obtain

$$\int_{\mathcal{K}_k^\dagger} \left( p_t(x) - p_t^{\top,k}(x) \right) p_t(x) dx \leq \frac{b\|p_t\|_\infty}{k^\gamma}. \quad (51)$$

The third term is trivial. Since  $p_t^{\top,k}(x) = 0$  for all  $x \in \mathcal{K}_k^\dagger$ , it follows that

$$\int_{\mathcal{K}_k^\dagger} \left( p_t^{\top,k}(x) - p_t(x) \right) p_t^{\top,k}(x) dx = 0. \quad (52)$$

Substituting Eqs. (50), (51) and (52) into Eq. (48) yields

$$\int \left( p_t^{\top,k}(x) - p_t(x) \right)^2 dx \leq \frac{\tilde{U}^\varepsilon}{k^{2-\varepsilon}} + \frac{b\|p_t\|_\infty}{k^\gamma} \leq \frac{U^\varepsilon}{k^{\min\{2-\varepsilon, \gamma\}}},$$

where  $U^\varepsilon = \tilde{U}^\varepsilon + b\|p_t\|_\infty$  and  $0 < \varepsilon < 2$ .

□

**Remark 12** A classical asymptotic approximation of the MISE (AMISE) for kernel density estimators built from i.i.d. samples is (see, e.g., [22] and note that we restrict ourselves to diagonal bandwidth matrices)

$$AMISE \equiv h^4 c(\phi, p_o) + \frac{c(\phi)}{Nh^{d_x}}, \quad (53)$$

where  $h > 0$  is the bandwidth parameter,  $c(\phi, p_o) > 0$  is a constant that depends on the kernel  $\phi$  and the target density (denoted  $p_o$  here and assumed twice differentiable),  $c(\phi) > 0$  is another constant depending

on  $\phi$  alone and  $N$  is the number of samples. If we substitute  $h = 1/k$  and  $N = k^{2d_x+2}$ , as given by our analysis, into the expression above, then we find that the MISE converges asymptotically as  $\frac{\tilde{c}(\phi, p_o)}{k^4}$ , for some constant  $\tilde{c}(\phi, p_o) > 0$ . We note, however, that

- Eq. (53) is only an asymptotic approximation of the MISE, whereas Theorems 4 and 5 give actual upper bounds for the MISE and the ISE that are valid for every  $k$ ;
- the AMISE of Eq. (53) is derived under the assumption that a size  $N$  sample drawn from the density  $p_t$  is available [44], whereas Theorems 4 and 5 hold true for the smoothing of any random measure  $\pi_t^N$  that satisfies  $\|(f, \pi_t^N) - (f, \pi_t)\|_p \leq \frac{c\|f\|_\infty}{\sqrt{N}}$  for some constant  $c$  and  $f \in B(\mathbb{R}^{d_x})$ .

#### 4.4 A simple example

There are several possible choices for the kernel function  $\phi(x)$  that comply with assumptions A.1 and A.2. In particular, the standard multivariate Gaussian density with unit covariance,

$$\phi_G(x) = \frac{1}{(2\pi)^{\frac{d_x}{2}}} \exp \left\{ -\frac{1}{2} \sum_{j=1}^{d_x} x_j^2 \right\},$$

the  $d_x$ -dimensional Laplacian pdf,

$$\phi_L(x) = \left( \frac{1}{2b} \right)^{d_x} \exp \left\{ -\frac{1}{b} \sum_{j=1}^{d_x} |x_j| \right\},$$

where  $b = \sqrt{\frac{1}{2d_x}}$ , and the Epanechnikov kernel  $\phi_E(x)$  of Eq. (15) are densities with bounded second order moment.

It is also straightforward to check assumption A.4 for  $\alpha = \mathbf{0}$  and  $\alpha = \mathbf{1}$ . In particular, for  $\alpha = \mathbf{0}$ , it is apparent that  $\phi_G, \phi_L, \phi_E \in C_b(\mathbb{R}^{d_x})$ . For  $\alpha = \mathbf{1}$ , the partial derivatives of the Gaussian and Laplacian kernels yield

$$\begin{aligned} D^{\mathbf{1}}\phi_G(x) &= \frac{(-1)^{d_x}}{(2\pi)^{\frac{d_x}{2}}} \prod_{l=1}^{d_x} x_l \exp \left\{ -\frac{1}{2} \sum_{j=1}^{d_x} x_j^2 \right\} \quad \text{and} \\ D^{\mathbf{1}}\phi_L(x) &= \frac{(-1)^{n^+}}{2^{d_x} b^{2d_x}} \exp \left\{ -\frac{1}{b} \sum_{j=1}^{d_x} |x_j| \right\}, \quad x \neq 0, \end{aligned}$$

respectively, where  $n^+ = |\{l \in \{1, \dots, d_x\} : x_l > 0\}|$  is the number of positive elements of  $x \in \mathbb{R}^{d_x}$ . It is not hard to verify that  $D^{\mathbf{1}}\phi_G \in C_b(\mathbb{R}^{d_x})$ , while  $D^{\mathbf{1}}\phi_L \in B(\mathbb{R}^{d_x})$ . As for the Epanechnikov kernel, it is easy to show that  $D^{\mathbf{1}}\phi_E(x) = 0 \forall x \in \mathbb{R}^{d_x}$ .

In the sequel, we consider a simple example consisting in the approximation of a Gaussian filtering density using the Epanechnikov kernel.

**Example 1** Consider the state-space system

$$p_0(x_0) = N(x_0; 0, \mathcal{I}_2), \quad X_t = AX_{t-1} + U_t, \quad Y_t = BX_t + V_t, \quad t = 1, 2, \dots \quad (54)$$

where  $N(x_0; 0, \mathcal{I}_2)$  is the bivariate Gaussian pdf with mean 0 and  $2 \times 2$  identity covariance matrix,  $\mathcal{I}_2$ ; the matrices  $A, B \in \mathbb{R}^{2 \times 2}$  are

$$A = \begin{bmatrix} 0.50 & -0.35 \\ 0.39 & -0.45 \end{bmatrix}, \quad B = \begin{bmatrix} 0.50 & 0.30 \\ -0.80 & 0.20 \end{bmatrix},$$

and  $U_t, V_t, t = 1, 2, \dots$ , are sequences of independent and identically distributed  $2 \times 1$  Gaussian vectors with zero mean and covariance  $\mathcal{I}_2$ . For this class of (linear and Gaussian) models the filtering pdf  $p_t$ ,  $t \geq 1$ , can be computed exactly using the Kalman filter [32] and, therefore, we have a reference for comparison with the approximations  $p_t^k$  produced by the particle filter with  $N(k) = k^{2(d_x+1)} = k^6$  samples.

For the simulation, we generated two sequences,  $x_0, x_1, \dots, x_T$  and  $y_1, \dots, y_T$  for  $T = 50$ , according to the model (54). Then, using the fixed data  $y_{1:T}$ , we run a Kalman filter to compute the Gaussian pdf  $p_T(x) = N(x; \bar{x}_T, \Sigma_T)$  exactly, where  $\bar{x}_T$  and  $\Sigma_T$  are the posterior mean and covariance at time  $T$ , respectively. For the same sequence  $y_{1:T}$ , we run independent particle filters with various values of  $k$  and  $N(k) = k^6$  particles each.

Figure 1 shows plots of the approximations  $p_T^k(x)$  for  $k = 4, 7, 10$  (constructed using the Epanechnikov kernel,  $\phi_E$ ) and the true pdf  $p_T(x)$ . The plots are drawn from a regular grid of points in  $\mathbb{R}^2$ , namely

$$x \in G_T = \{(x_1, x_2) : x_1 = -2.916 + 0.2n, \quad x_2 = -3.536 + 0.2n, \quad n = 1, 2, \dots, 42\} \quad (55)$$

(the offsets  $-2.916$  and  $-3.536$  correspond to the true posterior mean of  $X_t$ ). We can see that there is an obvious error for small  $k$ , while for  $k = 10$  the difference between  $p_T(x)$  and its approximation is negligible.


## 5 Applications

We illustrate the use of the convergence results in Section 3 by addressing two application problems: the computation of maximum a posteriori (MAP) estimators and the approximation of functionals of the filtering density,  $p_t$ . All through this section, we implicitly assume that the standard conditions of Remark 9 are satisfied.


### 5.1 MAP estimation

We tackle the problem of approximating the maximum a posteriori (MAP) estimator of the r.v.  $X_t$ . In particular, we address the numerical search of elements of the set


$$S_t = \arg \max_{x \in \mathbb{R}^{d_x}} p_t(x), \quad (56)$$




(a)  $k = 4$  and  $N(k) = 4,096$



(b)  $k = 7$  and  $N(k) = 117,649$



(c)  $k = 10$  and  $N(k) = 1,000,000$



(d)  $p_T(x)$  (exact)

Figure 1: Plots (a)-(c) display the approximations of the filtering density produced by the particle filter,  $p_T^k(x)$ , with an increasing number of particles  $N(k) = k^6$ , and an Epanechnikov kernel,  $\phi_E$ . The true pdf,  $p_T(x)$ , is shown in Plot (d) for comparison. The plots correspond to the discrete grid  $G_T$  in Eq. (55).

where  $s \in S_t$  if, and only if,  $p_t(s) = \max_{x \in \mathbb{R}^{d_x}} p_t(x)$ . Note that this is a relevant problem since MAP estimates are often used, e.g., in signal processing and engineering applications (see, e.g., [24, 37, 23]), and the density  $p_t(x)$  cannot be analytically found in general.

Let

$$S_t^k = \arg \max_{x \in \mathbb{R}^{d_x}} p_t^k(x)$$

be the set of MAP estimates for the approximation density  $p_t^k(x)$  and note that  $\hat{x}_k \in S_t^k$  if, and only if,  $p_t^k(\hat{x}_k) = \max_{x \in \mathbb{R}^{d_x}} p_t^k(x)$ . We can build a sequence of approximate estimates, denoted  $\{\hat{x}_k\}_{k \geq 1}$ , by taking one element from each set  $S_t^k$ ,  $k = 1, 2, \dots$ , at time  $t$ . If  $S_t$  is nonempty, then any convergent subsequence of  $\{\hat{x}_k\}_{k \geq 1}$  yields an arbitrarily accurate approximation of a true MAP estimator  $s \in S_t$ , as stated below.

**Theorem 6** Assume that  $S_t \neq \emptyset$  and take any convergent subsequence of  $\{\hat{x}_k\}_{k \geq 1}$ , denoted  $\{\hat{x}_{k_i}\}_{i \geq 1}$ . Let  $\hat{x} = \lim_{i \rightarrow \infty} \hat{x}_{k_i}$  be the limit of such subsequence. If  $p_t \in C_b(\mathbb{R}^{d_x})$ , then  $p_t(\hat{x}) = \max_{x \in \mathbb{R}^{d_x}} p_t(x)$ . In particular, if  $p_t(x)$  has a unique maximum, then  $S_t$  is a singleton and  $\lim_{i \rightarrow \infty} \hat{x}_{k_i} = \arg \max_{x \in \mathbb{R}^{d_x}} p_t(x)$ .

**Proof:** We prove the theorem by contradiction. Specifically, assume that  $p_t(\hat{x}) < \max_{x \in \mathbb{R}^{d_x}} p_t(x)$ . Then, choose some  $s \in S_t$ , so that  $p_t(s) = \max_{x \in \mathbb{R}^{d_x}} p_t(x)$  and  $p_t(\hat{x}) < p_t(s)$  and let

$$\epsilon \triangleq \frac{p_t(s) - p_t(\hat{x})}{3} > 0. \quad (57)$$

Now, choose a compact subset  $\mathcal{K} \subset \mathbb{R}^{d_x}$  that contain  $s$ ,  $\{\hat{x}_{k_i}\}_{i \geq 1}$  and  $\hat{x}$ . From Remark 11,  $\lim_{k \rightarrow \infty} \sup_{x \in \mathcal{K}} |p_t^k(x) - p_t(x)| = 0$  a.s., hence there exists  $m$  such that for all  $k \geq m$

$$\sup_{x \in \mathcal{K}} |p_t^k(x) - p_t(x)| < \epsilon. \quad (58)$$

Moreover, since  $p_t(x)$  is continuous at every point  $x \in \mathcal{K}$ , we can choose an integer  $i_0$  such that for all  $i \geq i_0$  we obtain

$$|p_t(\hat{x}_{k_i}) - p_t(\hat{x})| < \epsilon. \quad (59)$$

Now, choose an index  $\ell$  such that  $\ell \geq i_0$  and  $\ell \geq m$ . Then, for every  $i, k_i > \ell$ , we have

$$\begin{aligned} p_t^{k_i}(\hat{x}_{k_i}) - p_t^{k_i}(s) &= \overbrace{p_t^{k_i}(\hat{x}_{k_i}) - p_t(\hat{x}_{k_i})}^{< \epsilon} + \overbrace{p_t(\hat{x}_{k_i}) - p_t(\hat{x})}^{< \epsilon} \\ &\quad + \overbrace{p_t(\hat{x}) - p_t(s)}^{-3\epsilon} + \overbrace{p_t(s) - p_t^{k_i}(s)}^{< \epsilon} < 0, \end{aligned} \quad (60)$$

where the first term on the right-hand side,  $p_t^{k_i}(\hat{x}_{k_i}) - p_t(\hat{x}_{k_i}) < \epsilon$ , follows from inequality (58), the second term,  $p_t(\hat{x}_{k_i}) - p_t(\hat{x}) < \epsilon$ , follows from inequality (59), the third term,  $p_t(\hat{x}) - p_t(s) = -3\epsilon$ , is due to the definition in (57) and for the fourth term,  $p_t(s) - p_t^{k_i}(s) < \epsilon$ , is obtained from the inequality (58). Therefore,  $\hat{x}_{k_i} \notin \arg \max_{x \in \mathbb{R}^{d_x}} p_t^{k_i}(x)$  and we arrive at a contradiction. Hence,  $p_t(\hat{x}) = \max_{x \in \mathbb{R}^{d_x}} p_t(x)$ .

□

**Remark 13** Note that the whole sequence  $\{\hat{x}_k\}$  may not converge to a MAP estimate since it may, e.g., alternate between different elements of  $S_t$ .

Many global optimization algorithms, such as simulated annealing [5, 28] or accelerated random search [1], rely only on the evaluation of the objective function and Theorem 6 justifies their use with the approximation  $p_t^k(x)$ . Many other optimization procedures are based on the evaluation of derivatives of the objective function. For example, we may want to use a gradient search method to find a local maximum of  $p_t(x)$ , i.e., to find a solution of the equation

$$\nabla_x p_t(x) = 0, \quad (61)$$

where  $x = (x_1, \dots, x_{d_x})$  and

$$\nabla_x p_t(x) = \begin{bmatrix} \frac{\partial p_t}{\partial x_1} \\ \vdots \\ \frac{\partial p_t}{\partial x_{d_x}} \end{bmatrix} (x) = \begin{bmatrix} D^{\alpha_1} p_t \\ \vdots \\ D^{\alpha_{d_x}} p_t \end{bmatrix} (x),$$

with  $\alpha_i = (0, \dots, \overbrace{1}^{i\text{-th}}, \dots, 0)$ . Let  $x^*$  be a solution of (61), i.e.,  $\nabla_x p_t(x^*) = 0$ . Under the assumptions of Theorem 1, for every  $\epsilon > 0$  there exists  $k_\epsilon$  such that,  $\forall k > k_\epsilon$ ,

$$-\epsilon < D^{\alpha_i} p_t^k(x^*) < \epsilon \quad \text{a.s.}$$

Therefore,

$$\|\nabla_x p_t^k(x^*)\| = \sqrt{\sum_{i=1}^{d_x} \left( D^{\alpha_i} p_t^k(x^*) \right)^2} < \epsilon \sqrt{d_x}, \quad \forall k > k_\epsilon,$$

and, since  $\epsilon$  can be chosen as small as we wish,


$$\lim_{k \rightarrow \infty} \|\nabla_x p_t^k(x^*)\| = 0 \quad \text{a.s.},$$

which justifies the application of a gradient search procedure using the approximation of the filtering pdf.

**Example 2** We illustrate the application of a gradient search procedure using the same example as in Section 4.4. In particular, we consider the approximation of the maximum of the Gaussian filtering pdf  $p_T(x)$ ,  $T = 50$ , using a steepest descent method. Given an approximation  $p_T^k(x)$  of the filtering density constructed with the Gaussian kernel  $\phi_G$ , we run the iterative algorithm

$$\hat{x}_T(i+1)^k = \hat{x}_T(i)^k + a \nabla_x p_T^k(x) \big|_{x=\hat{x}_T(i)^k}, \quad i = 0, 1, 2, \dots \quad (62)$$

with initial condition  $\hat{x}_T(0)^k = (-2, -2)^\top$  and step-size parameter  $a = 0.1$ . This procedure yields a sequence of approximations  $\hat{x}_T(1)^k, \dots, \hat{x}_T(i)^k, \dots$  of the MAP estimator  $\hat{x}_T$ . Since for the model of Eq.



(a) With  $k = 5$  and  $N(k) = 15,625$ . (b) With  $k = 9$  and  $N(k) = 531,441$ . (c) With the true posterior pdf,  $p_t$ .

Figure 2: Trajectories of the gradient search algorithms. Plot (a) shows the estimates produced by the gradient search algorithm of Eq. (62) superimposed over a contour representation of  $p_T^{k=5}(x)$ . Plot (b) displays the estimates and contour graph for  $p_T^{k=9}(x)$ . Plot (c) shows the estimates produced by the gradient search algorithm of Eq. (63) superimposed over a contour representation of  $p_T(x)$ , for comparison.

(54) *it is possible to obtain  $p_T(x)$  exactly, we have also run a steepest descent search over the true filtering pdf, namely,*

$$\hat{x}_T(i+1) = \hat{x}_T(i) + a \nabla_x p_T(x)|_{x=\hat{x}_T(i)}, \quad i = 0, 1, 2, \dots, \quad (63)$$

*that generates the estimates  $\hat{x}_T(1), \dots, \hat{x}_T(i), \dots$  for the same initial condition and step size.*

*The results, using the same sequence of observations as in Section 4.4, are shown in Figure 2. Specifically, Figures 2(a) and 2(b) show the trajectories described by the estimates  $\hat{x}_T(1)^k, \dots, \hat{x}_T(i)^k, \dots$  superimposed over the contour plots of the approximate pdf  $p_T^k(x)$  for  $k = 5$  and  $k = 9$ , respectively (and  $N(k) = k^6$ ). For comparison, Figure 2(c) depicts the sequence  $\hat{x}_T(1), \dots, \hat{x}_T(i), \dots$  obtained from the search over the true density  $p_T(x)$ , together with the corresponding contour plot. We observe that both the pdf's and the trajectories described by the search algorithms are very similar.*

*Figure 3 depicts the sequences of estimates  $\hat{x}_T(1), \dots, \hat{x}_T(800)$  and  $\hat{x}_T(1), \dots, \hat{x}_T(800)$  computed by the search algorithms with the approximate and true gradients,  $\nabla p_T^k(x)$  and  $\nabla p_T(x)$ , respectively, component wise. We observe that, for  $k = 9$ , the approximation is tight in both dimensions.*

Similar arguments can be given for other optimization methods based on the derivatives of the density  $p_t$ . A relevant example is the Newton method (see, e.g., [3]) as it serves to illustrate the approximation of second order derivatives of  $p_t(x)$ . The Newton iteration for the computation of a solution of Eq. (61) has the form

$$\hat{x}_T(i+1) = \hat{x}_T(i) - \Delta_x p_T(x) \nabla_x p_T(x)|_{x=\hat{x}_T(i)}, \quad (64)$$

where  $\Delta_x p_T(x)$  is the  $d_x \times d_x$  Hessian matrix of  $p_T(x)$ .

**Example 3** *Again, we use the same example as in Section 4.4 for illustration and search the maximum of the Gaussian filtering pdf  $p_T(x)$ ,  $T = 50$ , by way of the Newton method. Given an approximation  $p_T^k(x)$*




Figure 3: Trajectories of the gradient search algorithms (approximate, with  $k = 9$ , and exact) on the horizontal (upper plot) and vertical (lower plot) planes.

of the filtering density constructed with the Gaussian kernel  $\phi_G$ , we run the iterative algorithm

$$\hat{x}_T(i+1)^k = \hat{x}_T(i)^k - \Delta_x p_T^k(x) \nabla_x p_T^k(x) \big|_{x=\hat{x}_T(i)^k}, \quad i = 0, 1, 2, \dots, \quad (65)$$


where both the Hessian matrix and the gradient vector are computed from the approximate density  $p_T^k(x)$  and the initial condition is  $\hat{x}_T(0)^k = (0.4, 0.4)$ . This procedure yields a sequence of approximations  $\hat{x}_T(1)^k, \dots, \hat{x}_T(i)^k, \dots$  of the MAP estimator  $\hat{x}_T$ . We have also run the Newton iteration over the true filtering pdf, i.e., the algorithm in (64), that generates the estimates  $\hat{x}_T(1), \dots, \hat{x}_T(i), \dots$  for the same initial condition.

The results for the fixed observation sequence  $y_{1:T}$  of Section 4.4 and Example 2, are shown in Figure 4. Specifically, Figure 4(a) shows the trajectory described by the estimates  $\hat{x}_T(1)^k, \dots, \hat{x}_T(i)^k, \dots$  superimposed over the contour plots of the approximate pdf  $p_T^k(x)$  for  $k = 7$  (and  $N(k) = k^6$ ). Figure 4(b) depicts the sequence  $\hat{x}_T(1), \dots, \hat{x}_T(i), \dots$  obtained from the search over the true density  $p_T(x)$  for reference.


Figure 5 displays the time evolution of the estimates  $\hat{x}_T(i)^k$  and  $\hat{x}_T(i)$ ,  $i = 1, \dots, 500$ , separately on each dimension. In general, the approximation of second order derivatives is less accurate than that of  $p_T(x)$  and its first-order derivatives for the same value of  $k$ .

## 5.2 Functionals of $p_t$

The result of Theorem 3 allows us to construct (rigorous) approximations of functionals of the form  $(f \circ p_t, \pi_t)$ , where  $\circ$  denotes composition and  $f$  is a Lipschitz-continuous and bounded real function. In



(a) With  $k = 7$  and  $N(k) = 117,649$ .



(b) With the true posterior pdf,  $p_t$ .

Figure 4: Trajectories of the Newton algorithms. Plot (a) shows the estimates produced by the Newton search algorithm of Eq. (65) superimposed over a contour representation of  $p_T^k(x)$  for  $k = 7$ . Plot (b) shows the estimates produced by the Newton search algorithm of Eq. (64) superimposed over a contour representation of  $p_T(x)$ , for comparison.




Figure 5: Trajectories of the Newton search algorithms (approximate, with  $k = 7$ , and exact) on the horizontal (upper plot) and vertical (lower plot) planes.

order to provide rates for the convergence of the particle-kernel approximations  $(f \circ p_t^k, \pi_t^{N(k)})$ , we again work with the sequence of hypercubes  $\mathcal{K}_k = [-M_k, M_k] \times \cdots \times [-M_k, M_k] \subset \mathbb{R}^{d_x}$  where  $M_k = \frac{1}{2}k^{\frac{\beta}{d_x p}}$  and  $0 < \beta < 1$ ,  $p > 3$  are constants with respect to  $k$ . Specifically, we have the following result.

**Theorem 7** Choose any bounded, Lipschitz continuous function  $f$ , i.e.,  $f \in B(\mathbb{R})$  and  $\forall x, y \in \mathbb{R}$

$$|f(x) - f(y)| \leq c_f |x - y|,$$

for some finite constant  $c_f > 0$ . If  $p_t \in B(\mathbb{R}^{d_x})$  and  $\pi_t(\mathcal{K}_k^\dagger) \leq \frac{b}{2}k^{-\gamma}$  for some constants  $\gamma, b > 0$  then

$$\left| (f \circ p_t^k, \pi_t^{N(k)}) - (f \circ p_t, \pi_t) \right| \leq \frac{Q_f^\varepsilon}{k^{\min\{1-\varepsilon, \gamma\}}} \quad (66)$$

where  $0 < \varepsilon < 1$  is an arbitrarily small constant and  $Q_f^\varepsilon$  is an a.s. finite random variable independent of  $k$ . In particular,

$$\lim_{k \rightarrow \infty} \left| (f \circ p_t^k, \pi_t^{N(k)}) - (f \circ p_t, \pi_t) \right| = 0 \quad \text{a.s.}$$

**Proof:** Consider first the absolute difference

$$\begin{aligned} \left| (f \circ p_t^k, \pi_t) - (f \circ p_t, \pi_t) \right| &= \left| \int [(f \circ p_t^k)(x) - (f \circ p_t)(x)] p_t(x) dx \right| \\ &\leq \int |(f \circ p_t^k)(x) - (f \circ p_t)(x)| p_t(x) dx, \end{aligned} \quad (67)$$

where the inequality holds because  $p_t(x) \geq 0$ . Using the Lipschitz continuity of  $f$  in the integral of Eq. (67) yields

$$\begin{aligned} \left| (f \circ p_t^k, \pi_t) - (f \circ p_t, \pi_t) \right| &\leq c_f \int |p_t^k(x) - p_t(x)| p_t(x) dx \\ &\leq c_f \|p_t\|_\infty \int |p_t^k(x) - p_t(x)| dx, \end{aligned} \quad (68)$$

where the second inequality follows from the assumption  $p_t \in B(\mathbb{R}^{d_x})$  (hence  $\|p_t\|_\infty < \infty$ ). Eq. (68) together with Theorem 3 readily yields

$$\left| (f \circ p_t^k, \pi_t) - (f \circ p_t, \pi_t) \right| \leq \frac{c_f \|p_t\|_\infty Q^\varepsilon}{k^{\min\{1-\varepsilon, \gamma\}}} \quad (69)$$

where  $0 < \varepsilon < 1$  is a constant and  $Q^\varepsilon$  is an a.s. finite random variable.

As a second step, consider the difference  $\left| (f \circ p_t^k, \pi_t^{N(k)}) - (f \circ p_t^k, \pi_t) \right|$ . Since  $f \in B(\mathbb{R})$ , it follows that  $\|f \circ p_t^k\|_\infty \leq \|f\|_\infty$  independently of  $k$  and an application of Proposition 1 yields

$$E \left[ \left| (f \circ p_t^k, \pi_t^{N(k)}) - (f \circ p_t^k, \pi_t) \right|^q \right] \leq \frac{c_t^q \|f\|_\infty^q}{N(k)^{\frac{q}{2}}} \leq \frac{c_t^q \|f\|_\infty^q}{k^{q(2d_x+1)}},$$

where  $q \geq 1$  and the second inequality holds because  $N(k) \geq k^{2(d_x+1)+1}$ . Using Lemma 1 with  $c = c_t^q \|f\|_\infty^q$  and  $\nu = 0$  (note that  $q(2d_x+1) \geq 2$  for any  $q, d_x \geq 1$ ), we readily obtain the convergence rate for the absolute error, i.e.,

$$\left| (f \circ p_t^k, \pi_t^{N(k)}) - (f \circ p_t^k, \pi_t) \right| \leq \frac{U^\varepsilon}{k^{1-\varepsilon}} \quad (70)$$

where  $0 < \varepsilon < 1$  is an arbitrarily small constant and  $U^\varepsilon \geq 0$  is an a.s. finite random variable.

To conclude, consider the triangle inequality

$$\left| (f \circ p_t^k, \pi_t^{N(k)}) - (f \circ p_t, \pi_t) \right| \leq \left| (f \circ p_t^k, \pi_t^{N(k)}) - (f \circ p_t^k, \pi_t) \right| + \left| (f \circ p_t^k, \pi_t) - (f \circ p_t, \pi_t) \right|. \quad (71)$$

Substituting (69) and (70) into (71) yields

$$\left| (f \circ p_t^k, \pi_t^{N(k)}) - (f \circ p_t, \pi_t) \right| \leq \frac{U^\varepsilon}{k^{1-\varepsilon}} + \frac{c_f \|p_t\|_\infty Q^\varepsilon}{k^{\min\{1-\varepsilon, \gamma\}}} \leq \frac{Q_f^\varepsilon}{k^{\min\{1-\varepsilon, \gamma\}}}, \quad (72)$$

where the random variable  $Q_f^\varepsilon = U^\varepsilon + c_f \|p_t\|_\infty Q^\varepsilon \geq 0$  is a.s. finite and independent of  $k$ .

□

In statistical signal processing and machine learning applications it is often of interest to evaluate the Shannon entropy of a probability measure  $\pi$ . Assuming that  $\pi$  has a density  $p$  w.r.t. the Lebesgue measure, the entropy of the probability distribution is

$$\mathcal{H}(\pi) = -(\log p, \pi) = - \int_{\mathcal{S}} \log[p(x)] p(x) dx,$$

where  $\mathcal{S}$  is the support of  $p$ . In the case of the filtering measure  $\pi_t$ , it is natural to think of a particle approximation of the entropy  $\mathcal{H}(\pi_t)$  constructed as

$$\mathcal{H}(\pi_t)^k = -(\log p_t^k, \pi_t^{N(k)}) = -\frac{1}{N(k)} \sum_{n=1}^{N(k)} \log p_t^k(x_t^{(n)}).$$

Unfortunately, the log function is neither bounded nor Lipschitz continuous and, therefore, Theorem 7 does not guarantee the convergence  $\mathcal{H}(\pi_t)^k \rightarrow \mathcal{H}(\pi_t)$ . Such a result, however, can be obtained, with a more specific argument, if we assume the support of the density  $p_t$  to be compact.

**Theorem 8** *Let the sequence of observations  $Y_{1:T} = y_{1:T}$  (for some large but finite  $T$ ) be fixed and assume that  $g_t^{y_t} \in B(\mathbb{R}^{d_x})$ ,  $(g_t^{y_t}, \xi_t) > 0$ ,  $1 \leq t \leq T$ , and the integrability conditions in Proposition 1, part (b), are satisfied. If there exists a compact set  $\mathcal{S} \subset \mathbb{R}^{d_x}$  such that  $\int_{\mathcal{S}} p_t(x) dx = 1$  and  $\inf_{x \in \mathcal{S}} p_t(x) > 0$ , then*

$$\lim_{k \rightarrow \infty} |\mathcal{H}(\pi_t)^k - \mathcal{H}(\pi_t)| = 0 \quad \text{a.s.}$$

**Proof:** We apply the triangle inequality to obtain

$$\begin{aligned} \left| (-\log p_t^k, \pi_t^{N(k)}) - (-\log p_t, \pi_t) \right| &\leq \left| (-\log p_t^k, \pi_t^{N(k)}) - (-\log p_t, \pi_t^{N(k)}) \right| \\ &\quad + \left| (-\log p_t, \pi_t^{N(k)}) - (-\log p_t, \pi_t) \right| \end{aligned} \quad (73)$$

and then analyze the two terms in the right-hand side of (73).

The first one can be expanded to yield

$$\begin{aligned} \left| (-\log p_t^k, \pi_t^{N(k)}) - (-\log p_t, \pi_t^{N(k)}) \right| &= \left| \frac{1}{N(k)} \sum_{i=1}^{N(k)} \log \frac{p_t(x_t^{(i)})}{p_t^k(x_t^{(i)})} \right| \\ &\leq \frac{1}{N(k)} \sum_{i=1}^{N(k)} \left| \log \frac{p_t(x_t^{(i)})}{p_t^k(x_t^{(i)})} \right|. \end{aligned} \quad (74)$$

The logarithm of a ratio  $x/y$  can be upper bounded as

$$\log \frac{x}{y} \leq \frac{\max\{x, y\}}{\min\{x, y\}} - 1, \quad (75)$$

hence applying (75) into (74) we arrive at

$$\left| (-\log p_t^k, \pi_t^{N(k)}) - (-\log p_t, \pi_t^{N(k)}) \right| \leq \frac{1}{N(k)} \sum_{i=1}^{N(k)} \left| \frac{\max\{p_t^k(x_t^{(i)}), p_t(x_t^{(i)})\}}{\min\{p_t^k(x_t^{(i)}), p_t(x_t^{(i)})\}} - 1 \right|. \quad (76)$$

However, from Theorem 2 and Remark 11,  $\lim_{k \rightarrow \infty} p_t^k(x)/p_t(x) = \lim_{N \rightarrow \infty} p_t(x)/p_t^k(x) = 1$  a.s. for every  $x \in \mathcal{S}$ . Moreover, since we have assumed  $\inf_{x \in \mathcal{S}} p_t(x) > 0$ , it follows that for any  $\epsilon > 0$  there exists  $k_\epsilon$  independent of  $x$  such that, for all  $k > k_\epsilon$ ,

$$\frac{\max\{p_t^k(x_t^{(i)}), p_t(x_t^{(i)})\}}{\min\{p_t^k(x_t^{(i)}), p_t(x_t^{(i)})\}} \leq 1 + \epsilon. \quad (77)$$

Substituting (77) into (76) yields, for all  $k > k_\epsilon$ ,

$$\left| (-\log p_t^k, \pi_t^{N(k)}) - (-\log p_t, \pi_t^{N(k)}) \right| \leq \epsilon \quad \text{a.s.}$$

Since  $\epsilon$  can be as small as we wish,

$$\lim_{k \rightarrow \infty} \left| (-\log p_t^k, \pi_t^{N(k)}) - (-\log p_t, \pi_t^{N(k)}) \right| = 0 \quad \text{a.s.} \quad (78)$$

The second term in (73) converges to 0 because of Proposition 1, part (b), i.e.,

$$\lim_{k \rightarrow \infty} \left| (-\log p_t, \pi_t^{N(k)}) - (-\log p_t, \pi_t) \right| = 0 \quad \text{a.s.} \quad (79)$$

and taking together Eqs. (78), (79) and (73) we arrive at

$$\lim_{k \rightarrow \infty} \left| (-\log p_t^k, \pi_t^{N(k)}) - (-\log p_t, \pi_t) \right| = 0 \quad \text{a.s.}$$

□

**Example 4** We continue to use the model of Section 4.4 to numerically illustrate the particle approximation of  $\mathcal{H}(\pi_t)$ . Since the densities  $p_t$  for this example are Gaussian with known covariance matrices  $\Sigma_t$ ,  $t = 1, 2, \dots$ , we can compute their associated Shannon entropies exactly, namely

$$\mathcal{H}(\pi_t) = \frac{1}{2} \log((2\pi e)^{d_x} |\Sigma_t|),$$

where  $|\Sigma_t|$  is the determinant of matrix  $\Sigma_t$ . Taking  $t = T = 50$  and using the same sequence of observations  $y_{1:50}$  as in Section 4.4, the resulting entropy is  $\mathcal{H}(\pi_T) = 2.5998$  nats.

Let us point out that, obviously, the Gaussian distribution has an infinite support and, therefore, the convergence result of Theorem 8 cannot be rigorously applied. However, the Gaussian pdf is light-tailed

|      | $k = 3$ | $k = 4$ | $k = 5$ |
|------|---------|---------|---------|
| mean | 0.0616  | 0.0370  | 0.0128  |
| std  | 0.0453  | 0.0249  | 0.0091  |

Table 1: Empirical mean and standard deviation of the entropy-approximation error,  $|\mathcal{H}(\pi_T) - \mathcal{H}(\pi_T)^k|$ , averaged over 30 independent simulations. The entropies are evaluated in nats.

and, as can be observed from Figure 1(d), it can be truncated within a compact (rectangular) support and still yield a faithful representation of the original distribution.

Table 1 displays the empirical mean and standard deviation of the absolute error

$$|\mathcal{H}(\pi_T) - \mathcal{H}(\pi_T)^k|$$

obtained through computer simulations for  $N(k) = k^6$  and  $k = 3, 4, 5$ . To be specific, we carried out 30 independent simulation runs for each value of  $k$ . We observe how both the mean error and its standard deviation reduce quickly as  $k$  is increased.

## 6 Summary

We have addressed the approximation of the sequence of filtering pdf's of a Markov state-space model using a particle filter. The numerical technique is conceptually simple. We collect the  $N$  particles generated by the sequential Monte Carlo algorithm and approximate the desired density as the sum of  $N$  scaled kernel functions located at the particle positions. The main contribution of the paper is the analysis of the convergence of such particle-kernel approximations. In particular, we have first proved the point-wise convergence of the approximation of the filtering density and its derivatives as the number of particles is increased and the kernel bandwidth is correspondingly decreased. Explicit convergence rates are provided and they are sufficient to prove that the approximation errors vanish a.s. Under mild additional assumptions on the chosen kernel, it is possible to extend the latter result to prove that the approximation error converges uniformly on the support of the filtering density (rather than point-wise) and a.s. to 0. We have also found an explicit convergence rate for the supremum of the approximation error. The analysis establishes a connection between the complexity of the particle filter and the bandwidth of the kernel function used for estimating the filtering pdf. For a fixed number of particles  $N$ , this relationship yields an optimal value of the bandwidth.

The uniform approximation result has a number of applications. We have first exploited it to prove the convergence, in total variation distance, of the continuous measure generated by the estimated density toward the true filtering measure. In a similar vein, we have also shown that the MISE of the sequence of approximate densities converges (quadratically with the kernel bandwidth) toward 0 when the state space is compact. For a truncated version of the density approximation, the (random) ISE is also shown to converge a.s. toward 0 without assuming compactness of the support. Although the convergence rate

found for the ISE is only quadratic (versus fourth order for the asymptotic approximation of the MISE in classical kernel density estimation theory), one should be aware that all the results obtained in this paper remain valid whenever the density estimator is obtained by smoothing a discrete random measure  $\pi_t^N$  that is “good enough” to estimate integrals of bounded functions in such a way that the  $L_p$  norms of the approximation error converge as  $\frac{c}{\sqrt{N}}$  (in particular, we do not require to have samples from the target density  $p_t$ ). As a consequence, the results obtained here can be applied to, e.g., kernel density estimators built from importance samples as in [45], or to the analysis of bootstrapped estimators as considered in [27].

We have also proved that the maxima of the approximate filtering density converge a.s. toward the true ones. Therefore, MAP estimation of the state at time  $t$  can be carried out using gradient and Newton search methods on the approximate filtering pdf. We remark that it is sound to apply such methods on the approximate function, since we have proved convergence also for its derivatives. The last application we consider is the approximation of functionals of the filtering pdf. We provide a general result that guarantees the convergence of the particle-kernel approximations for general bounded and Lipschitz continuous functionals of the filtering density. Finally, we prove that it is also possible to use the proposed constructs to approximate the Shannon entropy of densities with a compact support. In order to arrive at this result, we have also proved the convergence of the particle filter approximations of integrals of unbounded test functions under very mild assumptions (essentially, the integrability of the function up to fourth order). This is a departure from most existing approaches, which assume bounded test functions.

The theoretical results are complemented with some numerical examples, that include the approximation of a Gaussian density, its maximization using gradient and Newton methods and the approximation of its Shannon entropy. The reason to carry out all examples for a Gaussian pdf is that we can compute it exactly using a Kalman filter and, therefore, we can have a proper reference to assess the performance of the particle-kernel methods. In the case of the Shannon entropy approximation, the Gaussian distribution does not comply with the compact support assumption. However, being a light-tailed distribution, our simulations show that accurate estimates can still be computed.

## 7 Acknowledgements

The work of D. C. was partially supported by the EPSRC Grant number EP/H0005500/1. J. M. acknowledges the support of the Ministry of Science and Innovation of Spain (program Consolider-Ingenio 2010 CSD2008-00010 COMONSENS and project DEIPRO TEC2009-14504-C02-01).

## A Proof of Proposition 1

Part (a) of Proposition 1 is a straightforward consequence of [39, Lemma 1], hence we focus here on part (b).

We need to introduce an auxiliary result first. Let  $\{\theta_i\}_{i=1,\dots,N}$  be a collection of zero-mean random variables, conditionally i.i.d. given a  $\sigma$ -algebra  $\mathcal{G}$ . In particular,  $E[\theta_i|\mathcal{G}] = 0$  for all  $i$  and, for any integrable function  $h$ ,

$$E[h(\theta_i)h(\theta_j)|\mathcal{G}] = E[h(\theta_i)|\mathcal{G}]E[h(\theta_j)|\mathcal{G}]$$

for any  $i \neq j$ . We have the following basic inequality

**Lemma 2** *If the zero-mean, i.i.d. variables  $\theta_i$ ,  $i = 1, \dots, N$ , are  $p$ -integrable then*

$$E \left[ \left( \frac{1}{N} \sum_{i=1}^N \theta_i \right)^p | \mathcal{G} \right] \leq \frac{c}{N^{\frac{p}{2}}} E[\theta_1^p | \mathcal{G}], \quad (80)$$

where  $c$  is a constant independent of  $N$ . In particular, if  $p > 2$ ,

$$\lim_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{i=1}^N \theta_i \right| = 0 \quad a.s. \quad (81)$$

**Proof:** The inequality (80) is a straightforward consequence of the Marcinkiewicz-Zygmund inequality [38] (see also [31, Theorem 3] or [4, Chapter 10]). For  $p > 2$ , a standard Borel-Cantelli argument yields Eq. (81).  $\square$

Lemma 2 enables us to prove the convergence of particle approximations  $(f, \pi_t^N)$  toward  $(f, \pi_t)$  when the function  $f$  is integrable w.r.t.  $\pi_t$  but not necessarily bounded. We prove this result by induction in  $t$ . For  $t = 0$ , the samples  $x_0^{(i)}$ ,  $i = 1, \dots, N$ , are i.i.d. with common distribution  $\pi_0$ , hence the approximation  $\pi_0^N$  is constructed as  $\pi_0^N(dx) = \frac{1}{N} \sum_{i=1}^N \delta_{x_0^{(i)}}(dx)$ . It is, therefore, apparent that

$$E[(f, \pi_0^N) - (f, \pi_0)] = 0$$

and a simple application of Lemma 2 (with  $\theta_i = f(x_0^{(i)}) - (f, \pi_0)$ ) yields

$$\lim_{N \rightarrow \infty} |(f, \pi_0^N) - (f, \pi_0)| = 0 \quad a.s.$$

Next, we consider the induction hypothesis

$$\lim_{N \rightarrow \infty} |(f, \pi_t^N) - (f, \pi_t)| = 0 \quad a.s. \quad (82)$$

and study the particle approximations at time  $t + 1$ .

We start with the convergence of integrals of the form  $(f, \xi_{t+1})$ , where  $\xi_{t+1} = \tau_{t+1}\pi_t$ , as defined in Eq. (7). The particle approximation of  $\xi_t$  has the form

$$\xi_t^N(dx) = \frac{1}{N} \sum_{n=1}^N \delta_{\bar{x}_t^{(n)}}(dx),$$

hence the integral  $(f, \xi_{t+1}^N)$  has conditional mean

$$E[(f, \xi_{t+1}^N) | \mathcal{F}_t] = (f, \tau_{t+1}\pi_t^N),$$

given the  $\sigma$ -algebra  $\mathcal{F}_t = \sigma \left( x_l^{(n)}, \bar{x}_s^{(n)} : 1 \leq n \leq N, 0 \leq l \leq t, 0 < s \leq t \right)$ . In particular, the random variables  $\theta_i = f(\bar{x}_{t+1}^{(i)}) - (f, \tau_{t+1} \pi_t^N)$  are centered and i.i.d., and Lemma 2 yields

$$E \left[ |(f, \xi_{t+1}^N) - (f, \tau_{t+1} \pi_t^N)|^p | \mathcal{F}_t \right] = E \left[ \left| \frac{1}{N} \sum_{i=1}^N \theta_i \right|^p | \mathcal{F}_t \right] \leq \frac{\tilde{c}_t E[\theta_1^p | \mathcal{F}_t]}{N^{\frac{p}{2}}}, \quad (83)$$

where  $\tilde{c}_t$  is a constant w.r.t.  $N$ . Since both  $f$  and  $\bar{f}_{t+1}$  are  $p$ -integrable, it follows that  $E[\theta_1^p | \mathcal{F}_t]$  is a finite and non-negative random variable. Taking unconditional expectations on both sides of (83), we arrive at

$$E \left[ |(f, \xi_{t+1}^N) - (f, \tau_{t+1} \pi_t^N)|^p \right] \leq \frac{\tilde{c}_t}{N^{\frac{p}{2}}}, \quad (84)$$

where  $\tilde{c}_t = \tilde{c}_t E[E[\theta_1^p | \mathcal{F}_t]]$  is a constant w.r.t.  $N$ . From (84) it is straightforward to prove (e.g., using Lemma 1 or a Borel-Cantelli argument with  $p > 2$ ) that

$$\lim_{N \rightarrow \infty} |(f, \xi_{t+1}^N) - (f, \tau_{t+1} \pi_t^N)| = 0 \quad \text{a.s.} \quad (85)$$

Moreover, given the function  $\bar{f}_{t+1}(x) = \int f(x') \tau_{t+1}(dx' | x)$ , we can write  $(f, \tau_{t+1} \pi_t^N) = (\bar{f}_{t+1}, \pi_t^N)$  and  $(f, \tau_{t+1}, \pi_t) = (\bar{f}_{t+1}, \pi_t)$  and, as a consequence,

$$|(f, \tau_{t+1} \pi_t^N) - (f, \tau_{t+1} \pi_t)| = |(\bar{f}_{t+1}, \pi_t^N) - (\bar{f}_{t+1}, \pi_t)|.$$

Note that  $\bar{f}_{t+1}$  is a real function on  $\mathbb{R}^{d_x}$  and we have assumed that  $(\bar{f}_{t+1}^p, \pi_t) < \infty$ . Therefore, for  $p > 2$ , we can apply the induction hypothesis (82) to arrive at

$$\lim_{N \rightarrow \infty} |(\bar{f}_{t+1}, \pi_t^N) - (\bar{f}_{t+1}, \pi_t)| = 0 \quad \text{a.s.} \quad (86)$$

It is now straightforward to prove the convergence of  $(f, \xi_{t+1}^N)$ . Specifically, from the triangle inequality we obtain

$$\lim_{N \rightarrow \infty} |(f, \xi_{t+1}^N) - (f, \xi_{t+1})| \leq \lim_{N \rightarrow \infty} |(f, \xi_{t+1}^N) - (f, \tau_{t+1} \pi_t^N)| + \lim_{N \rightarrow \infty} |(f, \tau_{t+1} \pi_t^N) - (f, \xi_{t+1})| \quad (87)$$

and, since  $\xi_{t+1} = \tau_{t+1} \pi_t$ , substituting (85) and (86) into (87) yields

$$\lim_{N \rightarrow \infty} |(f, \xi_{t+1}^N) - (f, \xi_{t+1})| = 0 \quad \text{a.s.} \quad (88)$$

In order to analyze the approximations  $(f, \pi_{t+1}^N)$ , we first consider the measure

$$\bar{\pi}_{t+1}^N(dx) = \sum_{n=1}^N w_{t+1}^{(n)} \delta_{\bar{x}_{t+1}^{(n)}}(dx), \quad \text{where} \quad w_{t+1}^{(n)} = \frac{g_{t+1}^{y_{t+1}}(\bar{x}_{t+1}^{(n)})}{\sum_{k=1}^N g_{t+1}^{y_{t+1}}(\bar{x}_{t+1}^{(k)})}$$

obtained immediately after the computation of the weights in the particle filter. Integrals w.r.t.  $\bar{\pi}_{t+1}^N$  can be written in terms of  $g_{t+1}^{y_{t+1}}$  and  $\xi_{t+1}^N$ , namely

$$(f, \bar{\pi}_{t+1}^N) = \frac{(f g_{t+1}^{y_{t+1}}, \xi_{t+1}^N)}{(g_{t+1}^{y_{t+1}}, \xi_{t+1}^N)}.$$

This is natural, though, since from the Bayes theorem we readily derive the same relationship between  $\pi_{t+1}$  and  $\xi_{t+1}$ ,

$$(f, \pi_{t+1}) = \frac{(fg_{t+1}^{y_{t+1}}, \xi_{t+1})}{(g_{t+1}^{y_{t+1}}, \xi_{t+1})},$$

and the difference  $(f, \bar{\pi}_{t+1}^N) - (f, \pi_{t+1})$  can, therefore, be written as

$$\begin{aligned} (f, \bar{\pi}_{t+1}^N) - (f, \pi_{t+1}) &= \frac{(fg_{t+1}^{y_{t+1}}, \xi_{t+1}^N)}{(g_{t+1}^{y_{t+1}}, \xi_{t+1}^N)} - \frac{(fg_{t+1}^{y_{t+1}}, \xi_{t+1})}{(g_{t+1}^{y_{t+1}}, \xi_{t+1})} \pm \frac{(fg_{t+1}^{y_{t+1}}, \xi_{t+1}^N)}{(g_{t+1}^{y_{t+1}}, \xi_{t+1})} \\ &= \frac{(fg_{t+1}^{y_{t+1}}, \xi_{t+1}^N) - (fg_{t+1}^{y_{t+1}}, \xi_{t+1})}{(g_{t+1}^{y_{t+1}}, \xi_{t+1})} \\ &\quad + \frac{(fg_{t+1}^{y_{t+1}}, \xi_{t+1}^N)}{(g_{t+1}^{y_{t+1}}, \xi_{t+1}^N)} \times \frac{(g_{t+1}^{y_{t+1}}, \xi_{t+1}) - (g_{t+1}^{y_{t+1}}, \xi_{t+1}^N)}{(g_{t+1}^{y_{t+1}}, \xi_{t+1})}. \end{aligned} \quad (89)$$

Note that, since  $g_{t+q}^{y_{t+1}} \in B(\mathbb{R}^{d_x})$ , Eq. (88) yields

$$\lim_{N \rightarrow \infty} |(g_{t+1}^{y_{t+1}}, \xi_{t+1}) - (g_{t+1}^{y_{t+1}}, \xi_{t+1}^N)| = 0 \quad \text{a.s.} \quad (90)$$

and, since the assumptions in the Lemma impose that

$$(g_{t+1}^{y_{t+1}}, \xi_{t+1}) > 0, \quad (91)$$

it follows that

$$\lim_{N \rightarrow \infty} (g_{t+1}^{y_{t+1}}, \xi_{t+1}^N) > 0 \quad \text{a.s.} \quad (92)$$

Also as a consequence of  $g_{t+1}^{y_{t+1}} \in B(\mathbb{R}^{d_x})$ , the function  $fg_{t+1}^{y_{t+1}}$  is  $p$ -integrable and Eq. (88) guarantees, for  $p > 2$ , that

$$\lim_{N \rightarrow \infty} |(fg_{t+1}^{y_{t+1}}, \xi_{t+1}) - (fg_{t+1}^{y_{t+1}}, \xi_{t+1}^N)| = 0 \quad \text{a.s.} \quad (93)$$

and, in particular

$$\lim_{N \rightarrow \infty} (fg_{t+1}^{y_{t+1}}, \xi_{t+1}^N) < \infty \quad \text{a.s.} \quad (94)$$

Taking Eqs. (91) and (92) together, we deduce that the denominator of (89) is a.s. positive (as  $N \rightarrow \infty$ ). Similarly, Eqs. (90), (93) and (94) show that the numerator of (89) converges to 0 a.s. as  $N \rightarrow \infty$ . Therefore,

$$\lim_{N \rightarrow \infty} |(f, \bar{\pi}_{t+1}^N) - (f, \pi_{t+1})| = 0 \quad \text{a.s.} \quad (95)$$

On the other hand, the difference  $|(f, \pi_{t+1}^N) - (f, \bar{\pi}_{t+1})|$  also converges to 0 a.s. Indeed, given the  $\sigma$ -algebra  $\bar{\mathcal{F}}_t = \sigma(x_r^{(n)}, \bar{x}_s^{(n)}; \quad n = 1, \dots, N, \quad 0 \leq r < t, \quad 0 < s \leq t)$ ,

$$E[f(x_{t+1}^{(i)}) | \bar{\mathcal{F}}_{t+1}] = (f, \bar{\pi}_{t+1}^N), \quad i = 1, \dots, N,$$

hence the random variables  $\theta_i = f(x_{t+1}^{(i)}) - (f, \bar{\pi}_{t+1}^N)$  are centered and conditionally i.i.d. given  $\bar{\mathcal{F}}_{t+1}$ . As a consequence, Lemma 2 yields

$$E \left[ \left| \frac{1}{N} \sum_{i=1}^N \theta_i \right|^p | \bar{\mathcal{F}}_{t+1} \right] \leq \frac{\tilde{c}_{t+1} E[\theta_1^p | \bar{\mathcal{F}}_{t+1}]}{N^{\frac{p}{2}}}, \quad (96)$$

where  $\tilde{c}_{t+1}$  is a constant and  $\frac{1}{N} \sum_{i=1}^N \theta_i = (f, \pi_{t+1}^N) - (f, \bar{\pi}_{t+1}^N)$ . If we take full expectations on both sides of (96), we obtain

$$E \left[ |(f, \pi_{t+1}^N) - (f, \bar{\pi}_{t+1}^N)|^p | \bar{\mathcal{F}}_{t+1} \right] \leq \frac{\tilde{c}_{t+1}}{N^{\frac{p}{2}}}, \quad (97)$$

where  $\tilde{c}_{t+1} = \tilde{c}_{t+1} E[E[\theta_1^p | \bar{\mathcal{F}}_{t+1}]]$  is a finite constant. For  $p > 2$ , from Eq. (97), we readily deduce that

$$\lim_{N \rightarrow \infty} |(f, \pi_{t+1}^N) - (f, \bar{\pi}_{t+1}^N)| = 0 \quad \text{a.s.}, \quad (98)$$

using either Lemma 1 or a standard Borel-Cantelli argument.

Finally, taking together (95), (98) and the triangle inequality

$$|(f, \pi_{t+1}^N) - (f, \pi_{t+1})| \leq |(f, \pi_{t+1}^N) - (f, \bar{\pi}_{t+1}^N)| + |(f, \bar{\pi}_{t+1}^N) - (f, \pi_{t+1})|,$$

we arrive at  $\lim_{N \rightarrow \infty} |(f, \pi_{t+1}^N) - (f, \pi_{t+1})| = 0 \quad \text{a.s.}$

## B Proof of Lemma 1

Choose a constant  $\beta$  such that  $\nu < \beta < p - 1$  and define the random variable

$$U^{\beta,p} = \sum_{m=1}^{\infty} m^{p-1-\beta} (\theta^m)^p.$$

The variable  $U^{\beta,p}$  is obviously non-negative and, additionally, it has a finite mean,  $E[U^{\beta,p}] < \infty$ . Indeed, from Fatou's lemma

$$E[U^{\beta,p}] \leq \sum_{m=1}^{\infty} m^{p-1-\beta} E[(\theta^m)^p] \leq c \sum_{m=1}^{\infty} m^{-1-\beta+\nu},$$

where the second inequality follows from Eq. (20). Since  $\beta - \nu > 0$ , it follows that  $\sum_{m=1}^{\infty} m^{-1-(\beta-\nu)} < \infty$ , hence  $E[U^{\beta,p}] < \infty$ .

We use the so-defined random variable  $U^{\beta,p}$  in order to determine the convergence rate of  $\theta^k$ . Obviously,  $k^{p-1-\beta} (\theta^k)^p \leq U^{\beta,p}$  and solving for  $\theta^k$  yields

$$\theta^k \leq \frac{(U^{\beta,p})^{\frac{1}{p}}}{k^{1-\frac{1+\beta}{p}}}.$$

If we define  $\varepsilon = \frac{1+\beta}{p}$  and  $U^\varepsilon = (U^{\beta,p})^{\frac{1}{p}}$ , then we obtain the inequality

$$\theta^k \leq \frac{U^\varepsilon}{k^{1-\varepsilon}}.$$

Since  $E[U^{\beta,p}] < \infty$ , it follows that  $E[(U^\varepsilon)^p] < \infty$ , hence  $U^\varepsilon$  is a.s. finite. Also, we recall that  $\nu < \beta < p - 1$ , therefore  $\frac{1+\nu}{p} < \varepsilon < 1$ .

## References

- [1] M. J. Appel, R. Labarre, and D. Radulovic. On accelerated random search. *SIAM Journal on Optimization*, 14(3):708–730, 2003.
- [2] A. Bain and D. Crisan. *Fundamentals of Stochastic Filtering*. Springer, 2008.
- [3] J. F. Bonnans and C. Lemaréchal. *Numerical optimization: theoretical and practical aspects*. Springer-Verlag New York Inc, 2006.
- [4] Y. S. Chow and H. Teicher. *Probability theory independence, interchangeability, martingales (third edition)*. Springer, 2003.
- [5] A. Corana, M. Marchesi, C. Martini, and S. Ridella. Minimizing multimodal functions of continuous variables with the “simulated annealing” algorithm. *ACM Transactions on Mathematical Software*, 13(3):262–280, September 1987.
- [6] T. M. Cover and J. A. Thomas. *Elements of Information Theory*. Wiley-Interscience, New York (USA), 1991.
- [7] D. Crisan. Particle filters - a theoretical perspective. In A. Doucet, N. de Freitas, and N. Gordon, editors, *Sequential Monte Carlo Methods in Practice*, chapter 2, pages 17–42. Springer, 2001.
- [8] D. Crisan, P. Del Moral, and T.J. Lyons. Non-linear filtering using branching and interacting particle systems. *Markov Processes Related Fields*, 5(3):293–319, 1999.
- [9] D. Crisan and A. Doucet. Convergence of sequential Monte Carlo methods. Technical Report Cambridge University (CUED/FINFENG/TR381), 2000.
- [10] D. Crisan and A. Doucet. A survey of convergence results on particle filtering. *IEEE Transactions Signal Processing*, 50(3):736–746, March 2002.
- [11] T. A. Dean, S. S. Singh, A. Jasra, and G. W. Peters. Parameter estimation for hidden Markov models with intractable likelihoods. *arXiv*, 1103.5399v1[math.ST], 2011.
- [12] P. Del Moral. Non-linear filtering: interacting particle solution. *Markov Processes and Related Fields*, 2:555–580, 1996.
- [13] P. Del Moral. Non-linear filtering using random particles. *Theory of Probability and its Applications*, 40(4):690–701, 1996.
- [14] P. Del Moral. *Feynman-Kac Formulae: Genealogical and Interacting Particle Systems with Applications*. Springer, 2004.
- [15] P. Del Moral, A. Doucet, and S. Singh. Uniform stability of a particle approximation of the optimal filter derivative. *arXiv*, 1106.2525v1[math.ST], 2011.

- [16] P. Del Moral and L. Miclo. Branching and interacting particle systems. Approximations of Feynman-Kac formulae with applications to non-linear filtering. *Lecture Notes in Mathematics*, pages 1–145, 2000.
- [17] L. Devroye and L. Györfi. *Nonparametric Density Estimation: The  $L_1$  View*. Wiley, 1985.
- [18] R. Douc, O. Cappé, and E. Moulines. Comparison of resampling schemes for particle filtering. In *Proceedings of the 4<sup>th</sup> International Symposium on Image and Signal Processing and Analysis*, pages 64–69, September 2005.
- [19] A. Doucet, N. de Freitas, and N. Gordon. An introduction to sequential Monte Carlo methods. In A. Doucet, N. de Freitas, and N. Gordon, editors, *Sequential Monte Carlo Methods in Practice*, chapter 1, pages 4–14. Springer, 2001.
- [20] A. Doucet, N. de Freitas, and N. Gordon, editors. *Sequential Monte Carlo Methods in Practice*. Springer, New York (USA), 2001.
- [21] A. Doucet, S. Godsill, and C. Andrieu. On sequential Monte Carlo Sampling methods for Bayesian filtering. *Statistics and Computing*, 10(3):197–208, 2000.
- [22] T. Duong and M. L. Hazelton. Cross-validation bandwidth matrices for multivariate kernel density estimation. *Scandinavian Journal of Statistics*, 32(3):485–506, 2005.
- [23] L. Frenkel and M. Feder. Recursive expectation-maximization (EM) algorithms for time-varying parameters with applications to multiple target tracking. *IEEE Transactions Signal Processing*, 47(2):306–320, February 1999.
- [24] J. L. Gauvain and C. H. Lee. Bayesian learning for hidden markov model with gaussian mixture state observation densities. *Speech Communication*, 11(2-3):205–213, 1992.
- [25] S. Godsill, A. Doucet, and M. West. Maximum a posteriori sequence estimation using Monte Carlo particle filters. *Annals of the Institute of Statistical Mathematics*, 53(1):82–96, March 2001.
- [26] N. Gordon, D. Salmond, and A. F. M. Smith. Novel approach to nonlinear and non-Gaussian Bayesian state estimation. *IEE Proceedings-F*, 140(2):107–113, 1993.
- [27] P. Hall and K. H. Kang. Bootstrapping nonparametric density estimators with empirically chosen bandwidths. *The Annals of Statistics*, 29(5):1443–1468, 2001.
- [28] A.-R. Hedar and M. Fukushima. Derivative-free filter simulated annealing method for constrained continuous global optimization. *Journal of Global Optimization*, 35:521–549, 2006.
- [29] K. Heine and D. Crisan. Uniform approximations of discrete-time filters. *Advances in Applied Probability*, 40(4):979–1001, 2008.
- [30] X.L. Hu, T.B. Schon, and L. Ljung. A basic convergence result for particle filtering. *IEEE Transactions on Signal Processing*, 56(4):1337–1348, 2008.

- [31] R. Ibragimov and Sh. Sharakhmetov. Analogues of Khintchine, Marcinkiewicz-Zygmund and Rosenthal inequalities for symmetric statistics. *Scandinavian Journal of Statistics*, 26(4):pp. 621–633, 1999.
- [32] R. E. Kalman. A new approach to linear filtering and prediction problems. *Journal of Basic Engineering*, 82:35–45, 1960.
- [33] G. Kitagawa. Monte Carlo filter and smoother for non-Gaussian nonlinear state-space models. *J. Comput. Graph. Statist.*, 1:1–25, 1996.
- [34] H. R. Künsch. Recursive Monte Carlo filters: Algorithms and theoretical bounds. *The Annals of Statistics*, 33(5):1983–2021, 2005.
- [35] F. LeGland and N. Oudjane. Stability and uniform approximation of nonlinear filters using the Hilbert metric and application to particle filters. *Annals of Applied Probability*, pages 144–187, 2004.
- [36] J. S. Liu and R. Chen. Sequential Monte Carlo methods for dynamic systems. *Journal of the American Statistical Association*, 93(443):1032–1044, September 1998.
- [37] A. Logothetis and V. Krishnamurthy. Expectation maximization algorithms for map estimation of jump markov linear systems. *IEEE Transactions on Signal Processing*, 47(8):2139–2156, 1999.
- [38] J. Marcinkiewicz and A. Zygmund. Quelques théorèmes sur les fonctions indépendantes’). *Fund. Math*, 29:60–90, 1937.
- [39] J. Míguez, D. Crisan, and P. M. Djurić. On the convergence of two sequential Monte Carlo methods for maximum a posteriori sequence estimation and stochastic global optimization. *Statistics and Computing*, DOI: 10.1007/s11222-011-9294-4, 2011.
- [40] C. Musso, N. Oudjane, and F. LeGland. Improving regularised particle filters. In A. Doucet, N. de Freitas, and N. Gordon, editors, *Sequential Monte Carlo Methods in Practice*, chapter 12, pages 247–272. Springer, 2001.
- [41] K. Najim, E. Ikonen, and P. Del Moral. Open-loop regulation and tracking control based on a genealogical decision tree. *Neural Computing and Applications*, 15:339–349, 2006.
- [42] B. W. Silverman. *Density Estimation for Statistics and Data Analysis*. Chapman & Hall/CRC, Boca Raton (FL, USA), 1986.
- [43] J. S. Simonoff. *Smoothing Methods in Statistics*. Springer-Verlag, New York, 1996.
- [44] M. P. Wand and M. C. Jones. *Kernel Smoothing*. Chapman-Hall/CRC, 1995.
- [45] M. West. Approximating posterior distributions by mixture. *Journal of the Royal Statistical Society B*, pages 409–422, 1993.